

# Unimodal wave trains and solitons in convex FPU chains

Michael Herrmann\*

January 23, 2009

## Abstract

We consider atomic chains with nearest neighbour interactions and study periodic and homoclinic travelling waves which are called wave trains and solitons, respectively. Our main result is a new existence proof which relies on the constrained maximisation of the potential energy and exploits the invariance properties of an improvement operator. The approach is restricted to convex interaction potentials but refines the standard results as it provides the existence of travelling waves with unimodal and even profile functions. Moreover, we discuss the numerical approximation and complete localization of wave trains, and show that wave trains converge to solitons when the periodicity length tends to infinity.

Keywords: *Fermi-Pasta-Ulam chain, lattice travelling waves, constrained optimisation, complete localisation*

MSC (2000): 37K60, 47J30, 70F45, 74J30

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Variational approach</b>	<b>4</b>
2.1	Travelling waves as eigenfunctions of nonlinear integral equations	4
2.2	Variational structure for wave trains and solitons	6
2.3	Some pieces of functional analysis	8
<b>3</b>	<b>Wave Trains</b>	<b>11</b>
3.1	Existence results	11
3.2	Complete localisation of wave trains	13
<b>4</b>	<b>Solitons</b>	<b>16</b>
4.1	Existence of solitons for genuinely super-quadratic $\mathcal{P}$	19
4.2	Criteria for super-quadratic $\mathcal{P}$	21
4.3	Criteria for genuinely super-quadratic $\mathcal{P}$	23
4.4	Solitons as limits of wave trains	24

## 1 Introduction

We consider infinite chains of identical atoms with unit mass that are coupled by nearest neighbour interactions. The dynamics in such chains is governed by Newton's equations

$$\ddot{x}_j(t) = \Phi'(x_{j+1}(t) - x_j(t)) - \Phi'(x_j(t) - x_{j-1}(t)), \quad (1)$$

where  $x_j(t)$  denotes the position of the  $j$ th atom at time  $t$ , and  $\Phi$  is the interaction potential. Restating (1) in terms of atomic distances  $r_j(t) = x_{j+1}(t) - x_j(t)$  and atomic velocities  $v_j(t) = \dot{x}_j(t)$  we find

$$\dot{r}_j(t) = v_{j+1}(t) - v_j(t), \quad \dot{v}_j(t) = \Phi'(r_j(t)) - \Phi'(r_{j-1}(t)). \quad (2)$$

---

\*University of Oxford, Mathematical Institute, Centre for Nonlinear PDE (OxPDE), 24-29 St Giles', Oxford OX1 3LB, United Kingdom, michael.herrmann@maths.ox.ac.uk.

In this article we allow for arbitrary convex interaction potentials  $\Phi$  and refer to (1) as Fermi-Pasta-Ulam (FPU) chain although the potential in the original paper [FPU55] was a quartic polynomial.

FPU chains can be viewed as simple toy models for crystals and solids and allow to study some essential properties of nonlinear elastic materials. Even though (1) is a strong simplification of a real material it obeys a very complex behaviour and currently we are far from a complete understanding of its dynamical properties.

During the last decades, a lot of research addressed the existence and properties of travelling waves in FPU chains because they can be viewed as elementary waves and provide a lot of insight into the energy transport in nonlinear media. Such travelling waves solve nonlinear advance-delay differential equations and it is a fundamental mathematical problem to characterise the solution set to those equations.

A further motivation for the study of travelling waves is related to atomistic Riemann problems and self-thermalisation of FPU chains: Starting with piecewise constant initial data for the atomic distances and velocities, solutions to (2) are self-similar on a macroscopic scale and involve *dispersive shocks*, that are fan-like structures with strong microscopic oscillations, see Figure 1. It is known from the theory of integrable systems and numerical simulations that the oscillations within a dispersive shock can be described by *modulated travelling waves*, compare [FV99, DH08] and references therein.




Figure 1: The atoms in a dispersive shock self-organize into modulated travelling waves. *Left.* Snapshots of atomic distances and velocities against the scaled particle index for different macroscopic times. *Right.* Corresponding family of wave trains with super-sonic soliton as ‘homoclinic limit’; the picture shows the density plots of six local distribution functions in the  $(r, v)$ -plane. Modulation theory predicts that each of these local distribution functions corresponds to a travelling wave whose parameters depend on macroscopic time and particle index.

**Travelling waves and main result** *Travelling waves* are exact solutions to (2) and satisfy the ansatz<sup>1</sup>

$$r_j(t) = R(\varphi), \quad v_j(t) = V(\varphi), \quad \varphi = kj + \omega t, \quad (3)$$

with phase  $\varphi$ , wave number  $k$ , (negative) frequency  $\omega$ , and profile functions  $R$  and  $V$ . Inserting (3) into (2) we obtain the nonlinear advance-delay differential equations

$$\omega \frac{d}{d\varphi} R = \nabla_k^+ V, \quad \omega \frac{d}{d\varphi} V = \nabla_k^- \Phi'(R), \quad (4)$$

where  $\nabla_k^+$  and  $\nabla_k^-$  denote the forward and backward difference operators with shift  $k$ , respectively. Depending on the properties of the profile functions we distinguish the following cases:

<sup>1</sup>The ansatz (3) is slightly more general than the usual one, which assumes that the atomic positions  $x_j$  depend on the phase variable  $\varphi$ . In fact, the ansatz  $x_j(t) = X(kj + \omega t)$  is *not* invariant under Galilean transformations  $x \rightsquigarrow x + v_0 t$ , whereas (3) has this property as it respects  $r \rightsquigarrow r, v \rightsquigarrow v + v_0$ .

1. *Wave trains or periodic waves*:  $R$  and  $V$  are periodic,
2. *Solitons or homoclinic waves*:  $R$  and  $V$  are localized over a constant background state,
3. *Fronts or heteroclinic waves*:  $R$  and  $V$  connect different constant background states,
4. *Oscillatory fronts*:  $R$  and  $V$  connect different asymptotic wave trains.

Notice that our usage of ‘soliton’ is quite sloppy: Localized travelling waves are sometimes called ‘solitary waves’, and ‘soliton’ then refers to a solitary waves that survives collisions with other such waves unchanged.

In this article we show the existence of wave trains and solitons with unimodal and even profile functions  $R$  and  $V$ , where *even* means as usual invariance under  $\varphi \rightsquigarrow -\varphi$ , and *unimodal* functions are monotone for both  $\varphi \leq 0$  and  $\varphi \geq 0$ . Our main result can be stated as follows.

**Theorem 1.1.** *Under natural regularity assumptions on the convex potential  $\Phi$  there exists a four-parameter family of wave trains, and if  $\Phi$  satisfies additionally some super-quadratic growth conditions then there exists also a three-parameter family of solitons. Moreover, the profile functions  $R$  and  $V$  for both families are unimodal and even.*

Close to our work are [FW94, SW97, FV99, PP00, SZ07] where the existence of travelling waves is likewise studied in a variational framework. Therefore we shall compare both our method (§2.2) and results (§4.3) with those presented therein. We also refer to the numerical study [DEFW93], to [FM03] for existence results in 2D lattices, and to [Ioo00] which proves the existence of small-amplitude travelling waves by means of center-manifold reduction. Moreover, the existence of fronts is studied in [HR08a], and [SZ08] concerns oscillatory fronts in FPU chains with bi-harmonic potentials.

All results presented below concern solely wave trains and solitons in FPU chains, but the method can also be applied to other Hamiltonian lattices with convex potential energy  $\mathcal{P}$  as for instance Klein-Gordon chains with convex on-site potential, see [Her09], and atomic chains with next-nearest neighbour interactions.

We emphasize that we are not able to provide uniqueness results for travelling waves. Uniqueness of relative equilibria in Hamiltonian lattices is a notoriously difficult problem, and almost nothing is known about it. The only available results concern either the near-sonic limit [FP99], or systems where the travelling wave equation can be solved explicitly. Examples are the Toda chain [Tod81], the discrete nonlinear Schrödinger equation [HLM94], the harmonic chain, and the hard-sphere model [DHM06].

**Overview on the proof and organisation of the paper** In a first step we reformulate the travelling wave equation (4) in terms of a normalised profile function  $W \in \mathbb{L}^2$ . More precisely, we show that (4) can be transformed into a nonlinear eigenvalue equation

$$\omega^2 W = \partial \mathcal{P}(W), \tag{5}$$

where  $\mathcal{P}(W)$  is the potential energy of a travelling wave. The profile function  $W$  has no physical meaning, but determines  $R$  and  $V$  via  $R(\varphi) = r_0 + \int_{\varphi}^{\varphi+k} W(\tilde{\varphi}) d\tilde{\varphi}$  and  $V(\varphi) = v_0 + \omega W(\varphi)$ , where  $r_0$  and  $v_0$  are suitable normalization constants.

Our approach relies on a combination of variational and dynamical concepts, and can be summarized as follows.

1. Equation (5) is the Euler-Lagrange equation for the optimisation problem  $\mathcal{P}(W) \rightarrow \max$  subjected to the constraint  $W \in \mathcal{B}_\gamma$ , where  $\gamma$  is a free parameter,  $\mathcal{B}_\gamma \subset \mathbb{L}^2$  denotes the ball of radius  $\sqrt{2\gamma}$ , and  $\omega^2$  is the Lagrangian multiplier.
2. There exists an improvement dynamics  $W \mapsto \mathcal{T}_\gamma[W]$  on  $\mathcal{B}_\gamma$  that increases the potential energy. Moreover, each stationary point of this dynamics solves (5), and vice versa.

3. There exist non-trivial cones  $S$  that are invariant under the improvement dynamics. Consequently, each maximiser of  $\mathcal{P}$  in  $\mathcal{S} \cap \mathcal{B}_\gamma$  is a travelling wave, see Theorem 2.3.

We emphasize that the convexity of  $\Phi$  is essential for this approach as it is intimately related to both the properties of  $\mathcal{T}_\gamma$  and the existence of non-trivial invariant cones.

A major part of the mathematical analysis done in this paper is needed to show that there exists a maximizer of  $\mathcal{P}$  in  $\mathcal{S}_\gamma$ . In the wave-train setting we can use rather simple compactness arguments as the functional  $\mathcal{P}$  is continuous with respect to the weak topology in  $\mathbf{L}^2$ . In the soliton setting, however, we lack the weak compactness of  $\mathcal{P}$ , and the existence proof for maximizers requires more sophisticated arguments. Our main technical result in this context is Lemma 4.7, which implies that (for certain  $S$ ) the maximising sequences for  $\mathcal{P}$  in  $\mathcal{S}_\gamma$  are *localized*, and hence pre-compact in the strong topology.

This paper is organised as follows. In §2.1 we derive the fixed point equation (5) for both wave trains and solitons. The details of our variational approach are presented in §2.2, and §2.3 concerns the analytical properties of the underlying functionals and operators. In §3.1 we continue with the existence proof for wave trains and present some numerical simulations; the *complete localisation* of wave trains is studied in §3.2. The existence proof for solitons is contained in §4.1 and relies on a natural condition for the super-quadratic growth of the functional  $\mathcal{P}$ . In §4.2 and §4.3 we then discuss the corresponding properties for the interaction potential  $\Phi$ . Finally, inspired by the notion of  $\Gamma$ -convergence we show in §4.4 that wave trains converge to solitons when the periodicity length tends to infinity.

## 2 Variational approach

In this section we transform the travelling wave equation into a fixed point equation for a normalised profile function  $W$ , and describe our variational approach to existence results for both wave trains and solitons. To point out the key ideas we start with more formal consideration in §2.1 and §2.2, and postpone the analytical details to §2.3.

### 2.1 Travelling waves as eigenfunctions of nonlinear integral equations

In what follows we assume that the periodicity length of wave trains is given by  $2L$  with  $0 < L < \infty$ , and regard the corresponding profile functions  $R$  and  $V$  as defined on  $[-L, L]$ . Moreover, we identify the soliton case with  $L = \infty$  by considering  $R$  and  $V$  as functions on  $[-\infty, \infty]$ , or, equivalently, as functions on the Alexandrov compactification of  $\mathbb{R}$ . In other words, in both cases we impose the 'boundary conditions'  $R(L) = R(-L)$  and  $V(L) = V(-L)$ .

In what follows we denote by  $\mathbf{L}^2([-L, L])$  with  $L \in (0, \infty]$  the Lebesgue space of all square-integrable functions on  $[-L, L]$ , and if there is no risk of confusion we write  $\mathbf{L}^2$  instead of  $\mathbf{L}^2([-L, L])$ .

Our first aim is to transform the travelling wave equations for wave trains and solitons into eigenvalue equations for certain nonlinear integral operators defined on  $\mathbf{L}^2$ . For this purpose we define two linear averaging operators, and normalise the potential  $\Phi$ . More precisely, for given reference distance  $r_0$  we define the potential  $\Phi_{r_0}$  by

$$\Phi_{r_0}(r) = \Phi(r_0 + r) - \Phi(r_0) - \Phi'(r_0)r$$

which is *normalised* via  $\Phi_{r_0}(0) = \Phi'_{r_0}(0) = 0$  and  $\Phi''_{r_0}(0) = \Phi''(r_0)$ , and mention that this normalisation respects the convexity of  $\Phi$ . The eigenvalue equations for solitons and wave trains involve the averaging operators

$$(\bar{\mathcal{A}}_k W)(\varphi) = \int_{\varphi-k/2}^{\varphi+k/2} W(\tilde{\varphi}) d\tilde{\varphi},$$

and

$$\widehat{\mathcal{A}}_k W = \bar{\mathcal{A}}_k W - \frac{1}{2L} \int_{-L}^L \bar{\mathcal{A}}_k W(\tilde{\varphi}) d\tilde{\varphi} = \bar{\mathcal{A}}_k W - \frac{k}{2L} \int_{-L}^L W(\tilde{\varphi}) d\tilde{\varphi}.$$

Notice that the operator  $\bar{\mathcal{A}}_k$  is well defined and symmetric on  $\mathbb{L}^2$  for both finite and infinite  $L$ , compare Lemma 2.5 below, whereas  $\widehat{\mathcal{A}}_k$  is well defined for  $L < \infty$  only.

**Wave trains and normalisation via mean values** In order to reformulate the travelling wave equation (4) for  $L < \infty$  we introduce the *mean values* of wave trains by

$$r_{\text{av}} := \frac{1}{2L} \int_{-L}^L R(\varphi) d\varphi, \quad v_{\text{av}} := \frac{1}{2L} \int_{-L}^L V(\varphi) d\varphi.$$

**Observation 2.1.** *With the identification*

$$R(\varphi - k/2) = r_{\text{av}} + (\widehat{\mathcal{A}}_k W)(\varphi), \quad V(\varphi) = v_{\text{av}} + \omega W(\varphi), \quad (6)$$

for some profile function  $W$  with  $\frac{1}{2L} \int_{-L}^L W(\varphi) d\varphi = 0$  the integral equation

$$\omega^2 W = \widehat{\mathcal{A}}_k \Phi'_{r_{\text{av}}}(\widehat{\mathcal{A}}_k W). \quad (7)$$

is equivalent to the wave train equation (4).

*Proof.* First suppose that  $R$  and  $V$  solve (4), and let  $W = \omega^{-1}(V - v_{\text{av}})$ . Then, the first equation in (4) implies

$$\frac{d}{d\varphi} R(\varphi - k/2) = W(\varphi + k/2) - W(\varphi - k/2) = \frac{d}{d\varphi} \left( \widehat{\mathcal{A}}_k W \right)(\varphi),$$

and hence  $R(\varphi - k/2) = (\widehat{\mathcal{A}}_k W)(\varphi) + c_1$  for some constant  $c_1$ . Integrating this with respect to  $\varphi$  we find  $r_{\text{av}} = 0 + c_1$ , and hence (6)<sub>1</sub>. Moreover, the second equation in (4) provides

$$\begin{aligned} \omega^2 \frac{d}{d\varphi} W(\varphi) &= \Phi' \left( r_{\text{av}} + (\widehat{\mathcal{A}}_k W)(\varphi + k/2) \right) - \Phi' \left( r_{\text{av}} + (\widehat{\mathcal{A}}_k W)(\varphi - k/2) \right) \\ &= \Phi'_{r_{\text{av}}} \left( (\widehat{\mathcal{A}}_k W)(\varphi + k/2) \right) - \Phi'_{r_{\text{av}}} \left( (\widehat{\mathcal{A}}_k W)(\varphi - k/2) \right), \end{aligned}$$

and integration with respect to  $\varphi$  gives  $\omega^2 W = \widehat{\mathcal{A}}_k \Phi'_{r_{\text{av}}}(\widehat{\mathcal{A}}_k W) + c_2$  for some constant  $c_2$ . The condition  $\frac{1}{2L} \int_{-L}^L W(\varphi) d\varphi = 0$  now implies  $c_2 = 0$  and hence (6)<sub>2</sub>. Now let a solution  $W$  of (7) be given, and define both  $R$  and  $V$  by (6). Then, the first equation from (4) holds by construction, and the second one follows from (7) by differentiation with respect to  $\varphi$ .  $\square$

**Solitons and normalisation via background states** For solitons it is much more convenient to base the normalisation on the *background states*

$$r_{\text{bg}} := \lim_{\varphi \rightarrow \pm\infty} R(\varphi), \quad v_{\text{bg}} := \lim_{\varphi \rightarrow \pm\infty} V(\varphi),$$

because then the normalised profiles are localised.

**Observation 2.2.** *With the identification*

$$R(\varphi - k/2) = r_{\text{bg}} + (\bar{\mathcal{A}}_k W)(\varphi), \quad V(\varphi) = v_{\text{bg}} + \omega W(\varphi), \quad (8)$$

for some profile function  $W$  with  $\lim_{\varphi \rightarrow \pm\infty} W(\varphi) = 0$  the integral equation

$$\omega^2 W = \bar{\mathcal{A}}_k \Phi'_{r_{\text{bg}}}(\bar{\mathcal{A}}_k W). \quad (9)$$

is equivalent to the soliton equation (4).

*Proof.* Similar to that of Observation 2.1.  $\square$

We can consider the integral equation (9) also for  $L < \infty$ , and readily verify that each solution defines a wave train via (8). This normalisation for wave trains seems to be artificial because for finite  $L$  we have no immediate interpretation of the parameter  $r_{\text{bg}}$ . However, in §4.4 we rely on this setting and show that the  $L$ -periodic solutions  $W_L$  to (9) converge to solitons as  $L \rightarrow \infty$ , see Corollary 4.21, and this implies  $R_L(\pm L) \rightarrow r_{\text{bg}}$ .

**Parameter dependence of travelling waves** The travelling wave equation (4) obeys a simple scaling symmetry. In fact, the solution set to (4) is invariant under

$$R(\varphi) \rightsquigarrow R(\lambda\varphi), \quad V(\varphi) \rightsquigarrow V(\lambda\varphi), \quad k \rightsquigarrow \lambda^{-1}k, \quad \omega \rightsquigarrow \lambda^{-1}\omega, \quad L \rightsquigarrow \lambda^{-1}L, \quad (10)$$

with arbitrary  $\lambda > 0$ , corresponding to

$$W(\varphi) \rightsquigarrow \lambda W(\lambda\varphi) \quad \text{with} \quad r_{\text{bg}} \rightsquigarrow r_{\text{bg}}, \quad v_{\text{bg}} \rightsquigarrow v_{\text{bg}}, \quad r_{\text{av}} \rightsquigarrow r_{\text{av}}, \quad v_{\text{av}} \rightsquigarrow v_{\text{av}}.$$

Up to this scaling there remain four independent parameters for wave trains. A natural choice for nonlinear potentials is to fix the length parameter  $L$ , and to regard  $r_{\text{av}}$ ,  $v_{\text{av}}$ ,  $k$ , and  $\omega$  as independent parameters, see [DHM06]. In this paper, however, we prefer to fix the wave number  $k$ , so that wave trains are parametrised by  $r_{\text{av}}$ ,  $v_{\text{av}}$ ,  $L$ , and the *phase speed*  $\sigma := \omega/k$ . Accordingly, we parametrize solitons by  $r_{\text{bg}}$ ,  $v_{\text{bg}}$  and  $\sigma$ .

Notice that the velocity parameters  $v_{\text{av}}$  and  $v_{\text{bg}}$  do not appear in the travelling wave equations due to the Galilean invariance of FPU chains. Both  $v_{\text{av}}$  and  $v_{\text{bg}}$  are thus trivial parameters, which however become important when studying *modulated travelling waves* for which the parameters vary on a macroscopic scale, compare [DHM06, DH08].

## 2.2 Variational structure for wave trains and solitons

In view of the scaling and reformulation results from §2.1 we start to simplify our setting. In what follows we consider an arbitrary convex and smooth potential  $\Phi$  normalised by  $\Phi(0) = \Phi'(0) = 0$ . Moreover, we restrict to  $k = 1$ , so that the frequency  $\omega$  equals the phase speed  $\sigma$ , and consider solely the following two averaging operators

$$(\bar{\mathcal{A}}W)(\varphi) = \int_{\varphi^{-1/2}}^{\varphi^{+1/2}} W(\tilde{\varphi}) \, d\tilde{\varphi}, \quad (\hat{\mathcal{A}}W)(\varphi) = (\bar{\mathcal{A}}W)(\varphi) - \frac{1}{2L} \int_{-L}^L W(\varphi) \, d\varphi. \quad (11)$$

Moreover, in order to point out the similarities between wave trains and solitons we refer to an abstract averaging operator  $\mathcal{A}$ , which equals either  $\bar{\mathcal{A}}$  or  $\hat{\mathcal{A}}$ , and consider the general travelling wave equation

$$\sigma^2 W = \mathcal{A}\Phi'(\mathcal{A}W), \quad (12)$$

with  $W \in L^2([-L, L])$ , where  $L$  might be finite or infinite. In what follows we call  $W$  a *travelling wave* if and only if there exist  $\sigma^2 > 0$  such that (12) is satisfied.

Starting point for the variational formulation is the observation that (12) is the Euler-Lagrange equation to the action functional

$$\mathcal{L}(\sigma^2, W) = \mathcal{K}(\sigma^2, W) - \mathcal{P}(W),$$

where

$$\mathcal{K}(\sigma^2, W) = \frac{1}{2} \sigma^2 \int_{-L}^L W(\varphi)^2 \, d\varphi \quad \text{and} \quad \mathcal{P} = \int_{-L}^L \Phi((\mathcal{A}W)(\varphi)) \, d\varphi \quad (13)$$

are the *kinetic* and *potential* energy, respectively.

The first rigorous existence result for wave trains and solitons under super-quadratic growth assumptions for  $\Phi$  was given by Friesecke and Wattis [FW94]. The key idea is (in our notations) to minimise the  $L^2$ -norm of  $W$  under the constraint of prescribed potential energy, where  $1/\sigma^2$  plays the role of a Lagrangian multiplier. The existence of corresponding minimisers was then established by means of Lions' concentration-compactness principle [Lio84].

Smets and Willem [SW97] prove the existence of solitons by showing that for super-quadratic  $\Phi$  the functional  $\mathcal{L}$  satisfies the assumptions of (a modified) Mountain Pass Theorem. Recently, these results could be improved by Schwetlick and Zimmer. In [SZ07] they require the super-quadratic growth to hold only asymptotically, so certain double-well potentials are admissible.

Similarly to [SW97], Pankov and Pflüger [PP00] apply the Mountain Pass Theorem to wave trains, and pass to the limit  $L \rightarrow \infty$  by means of concentration compactness. Moreover, they present a different existence proof for solitons which is based on the Nehari manifold of  $\mathcal{L}$ .

A new idea for convex potentials  $\Phi$  was introduced by Filip and Venakides [FV99] by proposing to maximise the potential energy  $\mathcal{P}$  under the convex constraint  $W \in \mathcal{B}_\gamma$  with

$$\mathcal{B}_\gamma = \{W \in L^2 : \frac{1}{2} \|W\|_2^2 \leq \gamma\}.$$

A first advantage of this approach is that  $L < \infty$  implies the functional  $\mathcal{P}$  to be continuous with respect to the weak topology in  $L^2$ , so that the existence of wave trains follows from elementary principles of infinite-dimensional convex analysis. Secondly, the improvement operator  $\mathcal{T}_\gamma$  appears naturally in this context, and gives rise to effective approximation schemes for wave trains, see [Her05, DH08] and the numerical simulations below.

Our method is also based on the constrained maximization of the potential energy but yields improved results as it exploits several invariance properties  $\mathcal{T}_\gamma$ .

### Constrained maximisation of the potential energy via the improvement operator

In view of (12) we formally define the *improvement* operator  $\mathcal{T}_\gamma$  by

$$\mathcal{T}_\gamma[W] := \frac{\sqrt{2\gamma}}{\|\partial\mathcal{P}[W]\|_2} \partial\mathcal{P}[W],$$

where the operator  $\partial\mathcal{P}$  is the Gâteaux derivative of  $\mathcal{P}$ , that means  $\partial\mathcal{P}[W] = \mathcal{A}\Phi'(\mathcal{A}W)$  for all  $W \in L^2$ . By construction, each fixed point  $W$  of  $\mathcal{T}_\gamma$  is a travelling wave with  $\frac{1}{2} \|W\|_2^2 = \gamma$ , and vice versa, where the speed is given by  $\sigma^2 = \|\partial\mathcal{P}[W]\|_2 / \sqrt{2\gamma}$ .

In the next Section §2.3 we exploit the convexity of  $\Phi$  and derive the following building blocks for the existence proof.

1.  $\partial\mathcal{P}$  respects the positive cone

$$\mathcal{U} := \{W \in L^2 : W(-\varphi_1) = W(\varphi_1) \text{ and } W(\varphi_1) \geq W(\varphi_2) \text{ for almost all } 0 \leq \varphi_1 \leq \varphi_2 \leq L\},$$

which consists of all functions on  $[-L, L]$  that are even and unimodal. Moreover, for  $\mathcal{A} = \bar{\mathcal{A}}$  the operator  $\partial\mathcal{P}$  also respects

$$\mathcal{N} := \{W \in L^2 : W(\varphi) \geq 0 \text{ for almost all } \varphi \in [-L, L]\},$$

which is the cone of all non-negative functions.

2.  $\mathcal{T}_\gamma$  is well defined on  $\mathcal{B}_\gamma \setminus \mathcal{M}$  and maps into  $\partial\mathcal{B}_\gamma \setminus \mathcal{M}$ , where

$$\mathcal{M} := \{W \in L^2 : \mathcal{P}(W) = 0\}$$

is the set of all global minimisers of  $\mathcal{P}$ .

3.  $\mathcal{T}_\gamma$  increases the potential energy, that means  $\mathcal{P}(\mathcal{T}_\gamma W) \geq \mathcal{P}(W)$  for all  $W \notin \mathcal{M}$ , where equality holds if and only if  $W = \mathcal{T}_\gamma[W]$ .

We are now able to describe the key principle that provides the existence of travelling waves.

**Theorem 2.3.** *Let  $\mathcal{S} \subset \mathbf{L}^2$  be some positive cone that is invariant under the action of the operator  $\partial\mathcal{P}$ . Then, the set  $\mathcal{S}_\gamma \setminus \mathcal{M}$  with  $\mathcal{S}_\gamma = \mathcal{S} \cap \mathcal{B}_\gamma$  is invariant under the action of  $\mathcal{T}_\gamma$ , and each proper maximiser of  $\mathcal{P}$  in  $\mathcal{S}_\gamma$  is a fixed point of  $\mathcal{T}_\gamma$ , and hence a travelling wave.*

*Proof.* The invariance of  $\mathcal{S}_\gamma$  is implied by the assumption on  $\mathcal{S}$  and the properties of  $\mathcal{T}_\gamma$ . Now let  $W$  be a proper maximiser. Then,  $\mathcal{P}(\mathcal{T}_\gamma[W]) = \mathcal{P}(W) > \min \mathcal{P}|_{\mathcal{S}_\gamma}$  implies both  $\mathcal{T}_\gamma[W] = W$  and  $W \notin \mathcal{M}$ , and we conclude that  $W$  is in fact a travelling wave with  $\sigma^2 > 0$ .  $\square$

In what follows the cone  $\mathcal{S}$  is given by either  $\mathcal{U}$  or  $\mathcal{U} \cap \mathcal{N}$ . Since these cones are not open in  $\mathbf{L}^2$  the fact that each maximiser must satisfy the Euler-Lagrange equation (12) with multiplier  $\sigma^2$  is not clear a priori but provided by the invariance of  $\mathcal{S}_\gamma$  under  $\mathcal{T}_\gamma$ .

Theorem 2.3 yields only a sufficient condition for the existence of travelling waves. In fact, to show that  $\mathcal{P}$  attains its maximum in  $\mathcal{S}_\gamma$  is not trivial at all (at least in the soliton case), and requires a better understanding of the energy landscape in  $\mathcal{S}_\gamma$ . In our analysis we follow the direct approach and show that maximising sequences for  $\mathcal{P}$  are compact in some appropriate topology in  $\mathbf{L}^2$ . More precisely, for wave trains we use weak compactness, whereas in the soliton setting we establish the strong compactness for maximising sequences.

## 2.3 Some pieces of functional analysis

Here we prove the aforementioned properties of the improvement operator  $\mathcal{T}_\gamma$ . To this end we rely on following standing assumptions on the potential  $\Phi$ .

**Assumption 2.4.** *For given  $\gamma > 0$  we assume that the interaction potential  $\Phi$  has the following properties on the interval  $[-\sqrt{2\gamma}, \sqrt{2\gamma}]$ .*

1. (Smoothness)  $\Phi$  is at least  $\mathcal{C}^2$ ,
2. (Convexity)  $\Phi'' \geq 0$ ,
3. (Normalisation)  $0 = \Phi(0) = \Phi'(0)$  and  $\Phi''(0) = \beta \geq 0$ ,
4. (Non-triviality)  $\Phi$  does not vanish identically.

The restriction to the interval  $[-\sqrt{2\gamma}, \sqrt{2\gamma}]$  is natural in our context, because  $W \in \mathcal{B}_\gamma$  implies  $\|\mathcal{A}W\|_\infty \leq \sqrt{2\gamma}$ , see Lemma 2.5. As a consequence of Assumption 2.4 we find

$$0 \leq \Phi(r) \leq \frac{1}{2}r^2(\beta + o(|r|)), \quad \Phi'(-|r|) \leq 0 \leq \Phi'(|r|) \quad (14)$$

for all  $r$  with  $|r| \leq \sqrt{2\gamma}$ . Moreover, the non-triviality condition implies  $\mathcal{M} = \{W : \mathcal{P}(W) = 0\}$  and  $\mathcal{B}_\gamma \setminus \mathcal{M} \neq \{0\}$ , and hence each maximiser of  $\mathcal{P}$  in  $\mathcal{B}_\gamma$  is proper.

Within this section, the parameter  $L$  can take arbitrary values in  $(0, \infty]$ , and  $\mathbf{L}^p$  and  $W^{1,p}$  with  $1 \leq p \leq \infty$  denote the usual Lebesgue and Sobolev spaces on  $[-L, L]$ , where

$$\langle W_1, W_2 \rangle = \int_{-L}^L W_1(\varphi)W_2(\varphi) \, d\varphi$$

gives the inner product in  $\mathbf{L}^2$ .



## Properties of the averaging operators $\bar{\mathcal{A}}$ and $\hat{\mathcal{A}}$

We summarize some elementary properties of the averaging operators that are used in the proofs below.

**Lemma 2.5.** *For any  $L$ , the operator  $\bar{\mathcal{A}}$  is well defined on  $\mathbb{L}^2$  and has the following properties:*

1.  $\bar{\mathcal{A}}$  maps into  $\mathbb{L}^2 \cap \mathbb{L}^\infty$  with  $\|\bar{\mathcal{A}}W\|_\infty \leq \|W\|_2$  and  $\|\bar{\mathcal{A}}W\|_2 \leq \|W\|_2$ .
2.  $\bar{\mathcal{A}}$  maps into  $W^{1,2}$  with  $(\bar{\mathcal{A}}W)'(\varphi) = W(\varphi + 1/2) - W(\varphi - 1/2)$ .
3.  $\bar{\mathcal{A}}$  is self-adjoint on  $\mathbb{L}^2$ .
4. If a sequence  $(W_n)_n$  converges weakly in  $\mathbb{L}^2$  to some limit  $W_\infty$ , then  $(\bar{\mathcal{A}}W_n)_n$  converges strongly in  $\mathbb{L}^2([-L, L])$  for each finite  $L < \infty$  with  $L \leq L$ . In particular, for  $L < \infty$  the image of each bounded set in  $\mathbb{L}^2$  under the operator  $\bar{\mathcal{A}}$  is pre-compact in  $\mathbb{L}^2$  with respect to the strong topology.
5. In the wave train case ( $L < \infty$ ) the operator  $\bar{\mathcal{A}}$  is compact. Moreover, the  $m$ th eigenvalue ( $m=0, 1, 2, \dots$ ) is given by

$$\varrho_m = \Theta\left(\frac{m\pi}{2L}\right), \quad \Theta(\varrho) := \varrho^{-1} \sin(\varrho),$$

and the corresponding eigenspace is spanned by  $\cos\left(\frac{m\pi}{L}\cdot\right)$  and  $\sin\left(\frac{m\pi}{L}\cdot\right)$

6. In the soliton case ( $L = \infty$ ) the operator  $\bar{\mathcal{A}}$  is not compact anymore, because it has continuous spectrum  $\text{spec}_{\mathbb{L}^2}\bar{\mathcal{A}} = \{\Theta(\varrho) : \varrho \in \mathbb{R}\}$ . In particular,  $\text{spec}_{\mathbb{L}^2}\bar{\mathcal{A}}^2 = [0, 1]$ .

Moreover, for  $L < \infty$  we have  $\hat{\mathcal{A}} : \mathbb{L}^2 \rightarrow \mathbb{L}^2 \cap \mathbb{L}^\infty$  with  $\|\hat{\mathcal{A}}W\|_\infty \leq \|W\|_2$  and  $\|\hat{\mathcal{A}}W\|_2 \leq \|W\|_2$ .

*Proof.* Definition (11) gives

$$\bar{\mathcal{A}}V(\varphi) = \int_{-L}^L \bar{\chi}(\varphi - s)V(s) ds = \bar{\chi}(\varphi - \cdot) * V \quad (15)$$

where  $\bar{\chi}$  abbreviates the indicator function of the interval  $[-1/2, 1/2]$ , and  $*$  denotes the convolution operator. Hölder's inequality provides  $|\bar{\mathcal{A}}V(\varphi)|^2 \leq \int_{\varphi-1/2}^{\varphi+1/2} V(s)^2 ds$ , and from this we readily derive the first assertion. The proofs of the second and the third claim are then straight forward. Now suppose that  $W_n \rightarrow W_\infty$  weakly in  $\mathbb{L}^2([-L, L])$ , so that  $\bar{\mathcal{A}}W_n \rightarrow \bar{\mathcal{A}}W_\infty$  point-wise thanks to (15), and this implies the strong  $\mathbb{L}^2$ -convergence on each finite interval  $[-L, L]$  due to the uniform  $\mathbb{L}^\infty$ -bounds.

Towards the spectral properties of  $\bar{\mathcal{A}}$  we study how  $\Theta$  acts on plane waves. A direct calculation shows that each plane wave  $E_k(\varphi) = \varphi \mapsto e^{ik\varphi}$  satisfies the eigenvalue equation

$$\bar{\mathcal{A}}E_k = \Theta(k/2)E_k$$

pointwise, and this implies the fourth and the fifth assertion. Finally, for  $L < \infty$  we have

$$\hat{\mathcal{A}}V(\varphi) = \hat{\chi}(\varphi - \cdot) * V, \quad \hat{\chi}(\varphi) = \bar{\chi}(\varphi) - \frac{1}{2L}.$$

This implies  $|\hat{\mathcal{A}}W(\varphi)|^2 \leq \|\hat{\chi}\|_\infty \|\bar{\chi}(\varphi - \cdot)W\|_2^2 = |\bar{\mathcal{A}}W(\varphi)|^2$  and in turn the desired properties of  $\hat{\mathcal{A}}$ .  $\square$

As a consequence of Definition (11) and Lemma 2.5 we easily find

$$\ker_{\mathbb{L}^2}\bar{\mathcal{A}} = \left\{ W \in \mathbb{L}^2 : W(\cdot) = W(\cdot + 1), \int_{-1/2}^{1/2} W(\varphi) d\varphi = 0 \right\}. \quad (16)$$

In particular, the kernel of  $\bar{\mathcal{A}}$  is trivial if either  $L$  is irrational, or  $L = \infty$ . Moreover, for  $L < \infty$  we have

$$\ker_{\mathbb{L}^2}\hat{\mathcal{A}} = \ker_{\mathbb{L}^2}\bar{\mathcal{A}} \oplus \text{span}\{1\}. \quad (17)$$

### Properties of the potential energy functional $\mathcal{P}$

We rely on Assumption 2.4 and use standard methods from convex analysis to prove some properties of  $\mathcal{P}$  and its derivative. All results are formulated in terms of the abstract averaging operator  $\mathcal{A}$  and hold both in the wave train and the soliton case.

**Lemma 2.6.** *The functional  $\mathcal{P}$  is well-defined, bounded, continuous and Gâteaux-differentiable on  $\mathcal{B}_\gamma$ , and its derivative  $\partial\mathcal{P} = \mathcal{A} \circ \partial\Phi \circ \mathcal{A}$  is a monotone operator and maps  $\mathcal{B}_\gamma$  continuously into  $\mathbb{L}^2$ . Moreover, for arbitrary  $W_1, W_2 \in \mathcal{B}_\gamma$  we have*

$$\mathcal{P}(W_2) - \mathcal{P}(W_1) \geq \frac{m}{2} \|\mathcal{A}W_2 - \mathcal{A}W_1\|_2^2 + \langle \partial\mathcal{P}[W_1], W_2 - W_1 \rangle. \quad (18)$$

and

$$\langle \partial\mathcal{P}[W_2] - \partial\mathcal{P}[W_1], W_2 - W_1 \rangle \geq m \|\mathcal{A}W_2(\varphi) - \mathcal{A}W_1(\varphi)\|_2^2 \quad (19)$$

where the monotonicity constant  $m$  is given by  $m = \inf_{|r| \leq \sqrt{2\gamma}} \Phi''(r) \geq \beta$ .

*Proof.* For all  $W$  in  $\mathcal{B}_\gamma$  we have  $\|\mathcal{A}W\|_\infty \leq \sqrt{2\gamma}$ , and Assumption 2.4 implies

$$|\Phi'(r)| \leq C|r|, \quad \Phi(r) \leq \frac{1}{2}Cr^2$$

for all  $r$  with  $|r| \leq \sqrt{2\gamma}$ , where  $C = \sup_{|r| \leq \sqrt{2\gamma}} |\Phi''(r)| \geq \beta$ . Consequently, we find

$$0 \leq \mathcal{P}(W) \leq \frac{1}{2}C \|\mathcal{A}W\|_2^2 \leq \gamma C, \quad \|\mathcal{A}\partial\Phi[\mathcal{A}W]\|_p \leq C^p \|W\|_p \leq C^p \|W\|_p,$$

and all assertions concerning the continuity and boundedness of both  $\mathcal{P}$  and  $\partial\mathcal{P}$  follow immediately. Now let  $W_1, W_2 \in \mathcal{B}_\gamma$  be fixed, and notice that the convexity inequality  $(\Phi'(r_2) - \Phi'(r_1))(r_2 - r_1) \geq m(r_2 - r_1)^2$  with  $r_i = W_i(\varphi)$  implies (19) by integration w.r.t.  $\varphi$ . To prove (18), let  $\eta \in [0, 1]$  and consider  $W(\eta) := (1 - \eta)W_1 + \eta W_2 \in \mathcal{B}_\gamma$  as well as

$$p(\eta) := \mathcal{P}(W(\eta)).$$

The function  $p$  is well-defined and differentiable with respect to  $\eta$ , and using (19) we find

$$\begin{aligned} \frac{d}{d\eta} p(\eta) &= \langle \partial\mathcal{P}[W(\eta)], W_2 - W_1 \rangle \\ &\geq \eta^{-1} \langle \partial\mathcal{P}[W(\eta)] - \partial\mathcal{P}[W_1], \eta W_2 - \eta W_1 \rangle + \langle \partial\mathcal{P}[W_1], W_2 - W_1 \rangle \\ &= \eta^{-1} \langle \partial\mathcal{P}[W(\eta)] - \partial\mathcal{P}[W_1], W(\eta) - W_1 \rangle + \langle \partial\mathcal{P}[W_1], W_2 - W_1 \rangle \\ &\geq \eta^{-1} m \|\mathcal{A}W(\eta) - \mathcal{A}W_1\|_2^2 + \langle \partial\mathcal{P}[W_1], W_2 - W_1 \rangle \\ &\geq \eta m \|\mathcal{A}W_2 - \mathcal{A}W_1\|_2^2 + \langle \partial\mathcal{P}[W_1], W_2 - W_1 \rangle. \end{aligned}$$

Finally, we integrate the last estimate from  $\eta = 0$  to  $\eta = 1$ , and this gives (18).  $\square$

The convexity of  $\mathcal{P}$  implies that each trivial travelling wave with  $\sigma^2 = 0$  must belong to  $\mathcal{M}$ , the set of all minimisers of  $\mathcal{P}$ .

**Remark 2.7.** *We have  $\partial\mathcal{P}[W] \neq 0$  for all  $W \in \mathcal{B}_\gamma \setminus \mathcal{M}$ .*

*Proof.* Assume there exists some  $W \in \mathcal{B}_\gamma \setminus \mathcal{M}$  with  $\partial\mathcal{P}[W] = 0$ . Then, (19) with  $W_2 = 0$  and  $W_1 = W$  provides  $\mathcal{A}W = 0$  and hence  $\mathcal{P}(W) = 0$ , which is a contradiction.  $\square$

Notice that for non-degenerate potentials  $\Phi$  with  $\Phi(r) > 0$  for all  $r \neq 0$  we have  $\mathcal{M} = \ker \mathcal{A}$ , where  $\ker \mathcal{A}$  is given in (16) and (17).

### Properties of the improvement operator $\mathcal{T}_\gamma$

First we show that the cones  $\mathcal{U}$  and  $\mathcal{N}$  are invariant under the action of both  $\bar{\mathcal{A}}$  and  $\Phi'$ . Here again the convexity of  $\Phi$  enters as it guarantees that  $\Phi'$  increases monotonically.

**Lemma 2.8.** *The cones  $\mathcal{U}$  and  $\mathcal{N}$  are convex, closed under weak and strong convergence in  $\mathbf{L}^2$ , and are invariant under the action of  $\partial\mathcal{P}$  and  $\bar{\mathcal{A}}$ . Moreover,  $\mathcal{U}$  is also invariant under the action of  $\hat{\mathcal{A}}$  (for  $L < \infty$ ).*

*Proof.* The only non-trivial assertion is the invariance of  $\mathcal{U}$  under  $\bar{\mathcal{A}}$ . To prove this, we fix  $W \in \mathcal{U}$  and consider  $Y = \frac{d}{d\varphi}\bar{\mathcal{A}}W$  with  $Y(\varphi) = W(\varphi + 1/2) - W(\varphi - 1/2)$  thanks to Lemma 2.5. This function is odd as  $W(\varphi) = W(-\varphi)$  implies

$$Y(-\varphi) = W(-\varphi + 1/2) - W(-\varphi - 1/2) = W(\varphi - 1/2) - W(\varphi + 1/2) = -Y(\varphi).$$

Hence it remains to show that  $Y(\varphi) \leq 0$  for all  $\varphi \geq 0$ , which is equivalent to

$$W(\varphi + 1/2) \leq W(\varphi - 1/2), \quad \varphi \geq 0.$$

For  $1/2 \leq \varphi \leq L - 1/2$  this estimate follows from  $0 \leq \varphi - 1/2 \leq \varphi + 1/2$ . Moreover, for  $0 \leq \varphi \leq 1/2$  it holds thanks to  $W(-\varphi + 1/2) = W(\varphi - 1/2)$  and  $0 \leq -\varphi + 1/2 \leq \varphi + 1/2$ , and for  $L - 1/2 \leq \varphi \leq L$  it is a consequence of  $W(\varphi + 1/2) = W(2L - \varphi - 1/2)$  and  $2L - \varphi - 1/2 \geq \varphi - 1/2$ .  $\square$

**Lemma 2.9.** *The operator  $\mathcal{T}_\gamma$  maps  $\mathcal{B}_\gamma \setminus \mathcal{M}$  continuously into  $\partial\mathcal{B}_\gamma \setminus \mathcal{M}$  and satisfies*

$$\mathcal{P}(\mathcal{T}_\gamma[W]) - \mathcal{P}(W) \geq \frac{1}{2}m \|\mathcal{A}\mathcal{T}_\gamma[W] - \mathcal{A}W\|_2^2 \quad (20)$$

for all  $W \in \mathcal{B}_\gamma \setminus \mathcal{M}$ . Moreover, the equality sign holds if and only if  $W$  is a fixed point of  $\mathcal{T}_\gamma$ .

*Proof.* Lemma 2.6 and Remark 2.7 imply that  $\mathcal{T}_\gamma$  is well defined and continuous on  $\mathcal{B}_\gamma \setminus \mathcal{M}$ . Moreover,  $\|\mathcal{T}_\gamma[W_2]\|_2 = \sqrt{2\gamma}$  holds by definition. Now let  $W_1 \in \mathcal{B}_\gamma \setminus \mathcal{M}$  be fixed, and set  $W_2 := \mathcal{T}_\gamma[W_1]$  and  $\sigma_2^2 := \|\partial\mathcal{P}[W_1]\|_2 / \sqrt{2\gamma} > 0$ . Hence,  $\sigma_2^2 W_2 = \partial\mathcal{P}[W_1]$ , and from (18) we infer that

$$\begin{aligned} \mathcal{P}(W_2) - \mathcal{P}(W_1) - \frac{1}{2}m \|\mathcal{A}W_2 - \mathcal{A}W_1\|_2^2 &\geq \sigma_2^{-2} \langle W_2, W_2 - W_1 \rangle \\ &\geq \sigma_2^{-2} (\|W_2\|_2^2 - \|W_2\|_2 \|W_1\|_2), \end{aligned}$$

which gives (20) due to  $\|W_1\|_2 \leq \|W_2\|_2 = \sqrt{2\gamma}$ . Moreover, we find an equality sign in the second estimate if and only if  $\|W_2\|_2^2 = \|W_1\|_2^2 = \langle W_1, W_2 \rangle$ , that means if and only if  $W_1 = W_2$ .  $\square$

With Lemma 2.9 we have derived all ingredients that we had used in the proof of Theorem 2.3.

## 3 Wave Trains

As a first application of Theorem 2.3 we establish the existence of wave trains in §3.1 and proceed with some comments on the numerical computation of wave trains. Afterwards we study the complete localisation of wave trains in §3.2.

### 3.1 Existence results

Our first existence result concerns wave trains that are renormalised via their mean values. This corresponds to  $L < \infty$ ,  $\mathcal{A} = \hat{\mathcal{A}}$ , and  $\mathcal{S} = \mathcal{U}$ .

**Theorem 3.1.** *For each  $L < \infty$  and  $\gamma > 0$  there exist a unimodal and even wave train  $W$  such that  $\frac{1}{2} \|W\|_2^2 = \gamma$  and  $\sigma^2 W = \hat{\mathcal{A}}\Phi'(\hat{\mathcal{A}}W)$  for some  $\sigma^2 > 0$ .*

*Proof.* With respect to the weak topology in  $\mathbf{L}^2$ , the functional  $\mathcal{P}$  is continuous and the set  $\mathcal{S}_\gamma = \mathcal{U} \cap \mathcal{B}_\gamma$  is compact. Hence there exist a maximiser  $W$ , which is moreover proper, that means  $\sup_{\mathcal{S}_\gamma} \mathcal{P} = \mathcal{P}(W) > 0 = \min_{\mathcal{P}|_{\mathcal{S}_\gamma}}$ , because  $\Phi$  is non-trivial. The desired result now follows from Theorem 2.3.  $\square$

In view of Observation 2.1 and the scaling (10) we infer that Theorem 3.1 implies the existence of a four-parameter family of solutions  $(R, V)$  to the original travelling wave equation (4) with fixed  $L$ . This family is parametrised by  $r_{\text{av}}, v_{\text{av}}, k$ , and  $\gamma$ , and for nonlinear potentials we can moreover expect (at least locally) that  $\gamma$  can be replaced by  $\omega$ .

Similar existence result for wave trains in convex FPU chains are proven in [FV99, DHM06], but provide only  $W \in \partial\mathcal{B}_\gamma$ . Our method improves these results as it establishes the existence of wave trains with the additional property  $W \in \mathcal{U}$ , which in turn implies  $R \in \mathcal{U}$  and  $\sigma V \in \mathcal{U}$ . This sheds light on some observations from [DH08, HR08b]: The traces of travelling waves found in the numerical simulations of initial value problems for (1) typically encircle convex sets in the  $(r, v)$ -plane, compare the right picture in Figure 1. In particular, these curves have exactly two extrema in both the  $r$ -direction and the  $v$ -direction, and hence they correspond to unimodal profile functions  $R$  and  $V$ .

We proceed with some remarks concerning the uniqueness of wave trains. The norm constraint  $\frac{1}{2} \|W\|_2^2 = \gamma$  alone is not sufficient for uniqueness as the travelling wave equation is invariant under shifts in  $\varphi$ . Moreover, since the set of all  $2\tilde{L}$ -periodic functions, with  $m\tilde{L} = L$  for some  $m \in \mathbb{N}$ , is invariant under the action of  $\mathcal{T}_\gamma$ , we can construct a whole family of wave trains satisfying the norm constraint.

From these considerations we conclude that any uniqueness result for wave trains must prescribe further properties of the profile function  $W$ . Motivated by numerical simulations we conjecture, that for each  $\gamma$  there exists exactly one travelling wave with  $W \in \mathcal{U}$  and  $\frac{1}{2} \|W\|_2^2 = \gamma$ , but we are not able to prove this conjecture.

Finally, we use the same arguments as in the proof of Theorem 3.1 to derive a similar existence result for wave trains in the setting  $L < \infty$ ,  $\mathcal{A} = \bar{\mathcal{A}}$ , and  $\mathcal{S} = \mathcal{U} \cap \mathcal{N}$ .

**Lemma 3.2.** *For each  $L < \infty$  and  $\gamma > 0$  there exist a wave train  $W \in \mathcal{U} \cap \mathcal{N} \cap \partial\mathcal{B}_\gamma$  such that  $\sigma^2 W = \bar{\mathcal{A}}\Phi'(\bar{\mathcal{A}}W)$  for some  $\sigma^2 > 0$ .*

**Numerical computation of wave trains** It is natural to use the improvement dynamics

$$W \in \mathcal{S}_\gamma \mapsto \mathcal{T}_\gamma[W] \in \mathcal{S}_\gamma \tag{21}$$

for the approximation of wave trains, and a corresponding discrete scheme is readily derived and implemented. It was proven in [Her05] that the orbits generated by (21) are compact in the strong  $L^2$  topology, but from a theoretical point of view this result remains unsatisfactory due to the lack of uniqueness. So it is neither clear that maximisers of  $\mathcal{P}$  in  $\mathcal{S}_\gamma$  are unique, nor that all fixed points of  $\mathcal{T}_\gamma$  are (global) maximisers.

In numerical simulations, however, we found (21) to have good properties. For a wide class of potentials we observed rapid convergence to a unique limit independent of the chosen initial data. In Figure 2 we present the numerically computed profiles  $W$  for different values of  $\gamma$  with  $\mathcal{A} = \bar{\mathcal{A}}$  and

$$\Phi(r) = \cosh(r) - 1. \tag{22}$$

For small  $\gamma$  we can approximate the potential by the harmonic one  $\Phi_{\text{harm}}(r) = \Phi''(0)r^2$ , and hence the profile  $W$  is close to a rescaled plane wave. For increasing  $\gamma$ , however, the nonlinearity dominates and the profile function becomes tighter. Figure 3 shows the corresponding *traces* in the  $(r, v)$ -plane, these are the curves

$$\varphi \mapsto (\bar{\mathcal{A}}W(\varphi), \sigma W(\varphi)) \cong (R(\varphi), V(\varphi)),$$

compare Observation (2.2). Surprisingly we find a nested family of curves, that mean the traces for different values of  $\gamma$  do not intersect but fill out a convex region. We are not able to prove this observation but mention that a similar phenomenon occurs when FPU chains generate dispersive shocks, see [DHR06, HR08b].




Figure 2: Profile functions  $W$  for several values of  $\gamma$  with  $L = 2$  and  $\Phi$  as in (22).




Figure 3: Traces for the wave trains from Figure 2.

### 3.2 Complete localisation of wave trains

It is well known for strongly nonlinear potentials that in certain limits the wave trains (and even the solitons) localize completely, in the sense that – under a suitable rescaling – the profile functions  $W$  converge to the indicator function of an interval plus a constant background state. Such profile functions are, up to renormalisation, equal to the profile functions of travelling waves in the hard-sphere model for the atomic chain, in which all atomic interactions are described by elastic collisions. Thus the effect of localisation can often be linked in a natural way to the high energy limit of travelling waves. For more details we refer to [Tod81, FM02, Her05, DHM06].

In this section we discuss the localisation phenomenon in our context, and aim to derive a localisation criterion for wave trains. To keep the presentation simple we solely consider non-negative and unimodal profile functions, that means we investigate the localisation of solutions to (12) with  $L < \infty$ ,  $\mathcal{A} = \bar{\mathcal{A}}$ , and  $\mathcal{S} = \mathcal{U} \cap \mathcal{N}$ . Moreover, for our purpose it is sufficient to assume that the localised limit profile is given by

$$W_{\text{CL}}(\varphi) = \chi_{[-\frac{1}{2}, \frac{1}{2}]}(\varphi) = \begin{cases} 1 & \text{if } |\varphi| \leq \frac{1}{2}, \\ 0 & \text{if } |\varphi| > \frac{1}{2}, \end{cases} \quad (23)$$

see Figure 4. It is easy to check that this profile satisfies

$$\|W_{\text{CL}}\|_2 = 1, \quad (\bar{\mathcal{A}}W_{\text{CL}})(\varphi) = \max\{1 - |\varphi|, 0\}, \quad \mathcal{P}(W_{\text{CL}}) = 2 \int_0^1 \Phi(s) ds. \quad (24)$$

In what follows we consider sequences  $(\Phi_n)_n$  of rescaled potentials, where each  $\Phi_n : [0, 1] \rightarrow \mathbb{R}$  satisfies Assumption 2.4. Moreover, we refer to a sequence of profile functions  $W_n \subset L^2$  as a *corresponding sequence*

of maximisers, if  $W_n$  is a maximiser of  $\mathcal{P}_n$  in  $\mathcal{S}_{1/2}$  for each  $n$ , where  $\mathcal{P}_n$  is the potential energy functional (13) corresponding to  $\Phi_n$ .




Figure 4: The functions  $W_{\text{CL}}$  and  $\bar{\mathcal{A}}W_{\text{CL}}$ .

We say, a sequence of such potentials  $(\Phi_n)_n$  has the *complete localisation property on  $[0, 1]$*  if any corresponding sequence of maximisers converges strongly in  $L^2$  to  $W_{\text{CL}}$ . Our main result in this section is a necessary condition for the complete localisation of wave trains and is implied by the following observation.

**Lemma 3.3.** *We have  $\|\bar{\mathcal{A}}W\|_\infty < \|\bar{\mathcal{A}}W_{\text{CL}}\|_\infty = 1$  for any  $W \in \mathcal{S}_{1/2}$  with  $W \neq W_{\text{CL}}$ .*

*Proof.*  $W \in \mathcal{U} \cap \mathcal{N}$  implies  $\|\bar{\mathcal{A}}W\|_\infty = (\bar{\mathcal{A}}W)(0)$  and Hölder's inequality provides

$$\begin{aligned} (\bar{\mathcal{A}}W)^2(0) &= \left( \int_{-1/2}^{1/2} W(\varphi) \, d\varphi \right)^2 = \left( \int_{-1/2}^{1/2} W_{\text{CL}}(\varphi)W(\varphi) \, d\varphi \right)^2 \\ &\leq \left( \int_{-1/2}^{1/2} W_{\text{CL}}(\varphi)^2 \, d\varphi \right) \left( \int_{-1/2}^{1/2} W(\varphi)^2 \, d\varphi \right) = \int_{-1/2}^{1/2} W(\varphi)^2 \, d\varphi \end{aligned} \quad (25)$$

$$\leq 1 = (\bar{\mathcal{A}}W_{\text{CL}})^2(0). \quad (26)$$

Moreover, the estimate in (25) is strict unless there exist a constant  $c$  such that

$$W(\varphi) = cW_{\text{CL}}(\varphi) \quad \text{for almost all } \varphi \in [-\tfrac{1}{2}, \tfrac{1}{2}], \quad (27)$$

whereas the estimate in (26) is strict except for

$$\int_{-1/2}^{1/2} W(\varphi)^2 \, d\varphi = 1. \quad (28)$$

Now suppose that both (27) and (28) are satisfied. Then we have  $c = 1$ , and the norm constraint  $\|W\|_2^2 \leq 1$  implies  $W(\varphi) = 0 = W_{\text{CL}}(\varphi)$  for almost all  $\varphi \notin [-\frac{1}{2}, \frac{1}{2}]$ , and hence  $W = W_{\text{CL}}$ .  $\square$

**Lemma 3.4.** *The sequence  $(\Phi_n)_n$  has the complete localisation property on  $[0, 1]$  provided that the following two conditions are satisfied.*

1.  $\mathcal{P}_n(W_{\text{CL}}) = 1$  for all  $n$ .
2.  $\Phi_n$  converges uniformly and essentially monotonically to 0 on each interval  $[0, r_0]$  with  $0 < r_0 < 1$ . That means for any  $r_0$  we have  $\sup_{0 \leq r \leq r_0} \Phi_n(r) \rightarrow 0$  as  $n \rightarrow \infty$ , and there exists  $n_0(r_0)$  such that

$$0 \leq \Phi_{n_2}(r) \leq \Phi_{n_1}(r)$$

for all  $n_2 > n_1 > n_0(r_0)$  and all  $0 \leq r \leq r_0$ .

*Proof.* First we assume additionally that  $W_n \rightarrow W_\infty$  weakly in  $L^2$ , and suppose for contradiction that  $W_\infty \neq W_{\text{CL}}$ . Then,  $\|\bar{A}W_\infty\|_\infty < 1$  thanks to Lemma 3.3, and since  $\bar{A}W_n \rightarrow \bar{A}W_\infty$  point-wise and  $\|\bar{A}W_n\|_\infty = (\bar{A}W_n)(0)$  we find some  $0 < r_0 < 1$  such that  $\|\bar{A}W_n\|_\infty \leq r_0$  for almost all  $n$ . Therefore,  $\mathcal{P}_{n_1}(W_{n_2}) \geq \mathcal{P}_{n_2}(W_{n_2})$  holds for all  $n_2 \geq n_1$  and all sufficiently large  $n_1$ , and since each  $W_n$  is a maximiser for  $\mathcal{P}_n$  we also have  $\mathcal{P}_{n_2}(W_{n_2}) \geq \mathcal{P}_{n_2}(W_{\text{CL}}) \geq 1$ . We conclude that  $\mathcal{P}_{n_1}(W_{n_2}) \geq 1$  and passing to the limits  $n_2 \rightarrow \infty$  and  $n_1 \rightarrow \infty$  we obtain  $\liminf_{n \rightarrow \infty} \mathcal{P}_n(W_\infty) \geq 1$ . However,  $\|\bar{A}W_\infty\|_\infty \leq r_0 < 1$  implies  $\lim_{n \rightarrow \infty} \mathcal{P}_n(W_\infty) = 0$ , the desired contradiction. The result obtained so far implies that  $W_{\text{CL}}$  is the unique accumulation point of a maximising sequence, and this yields the weak convergence to  $W_{\text{CL}}$  for any maximising sequence. Finally, the strong convergence follows from  $1 = \|W_{\text{CL}}\|_2 = \|W_n\|_2$  for all  $n$ .  $\square$

Notice that the two conditions from Lemma 3.4 imply  $\Phi_n(r) \rightarrow 0$  for all  $0 \leq r < 1$ , but  $\Phi_n(1) \rightarrow \infty$  as  $n \rightarrow \infty$ , and for this reason it is not clear whether or not the complete localisation (convergence of maximisers) implies the convergence of maxima. The simulations from Figure 6, however, provide evidence for  $\liminf_{n \rightarrow \infty} \mathcal{P}_n(W_n) > 1 = \lim_{n \rightarrow \infty} \mathcal{P}_n(W_{\text{CL}})$ .

Our first application concerns the maximiser for homogeneous potentials of large degree. For an illustration we refer to the numerical results in Figure 5 and 6.

**Example 3.5.** *The family of potentials  $\Phi_q(r) = \frac{q+1}{2}r^q$  with  $c > 0$  and  $q > 2$  has the complete localisation property on  $[0, 1]$  for  $q \rightarrow \infty$ .*

*Proof.*  $\mathcal{P}_q(W_{\text{CL}}) = 1$  follows from (24) by a direct computation, and for fixed  $0 < r_0 < 1$  we choose  $q_0$  such that  $1 + (q_0 + 1) \ln r_0 < 0$ . Then we find  $\partial_q(\frac{q+1}{2}r^q) = \frac{1}{2}r^q(1 + (q + 1) \ln r) < 0$  for all  $0 \leq r < r_0$  and  $q > q_0$ .  $\square$




Figure 5: Profile functions  $W$  with  $\gamma = \frac{1}{2}$ ,  $L = 2$ , and potentials  $\Phi_q$  as in Remark 3.5. This example describes the wave trains for homogeneous potentials in the limit of increasing degree.




Figure 6: Speed  $\sigma$  and potential energy  $\mathcal{P}(W)$  versus  $q$  for the wave trains from Figure 5.

The second candidate for the complete localisation property is related to the limit  $\gamma \rightarrow \infty$  for fixed potential  $\Phi$ . For this purpose we consider the rescaled potentials

$$\Phi_\gamma(r) := \frac{\Phi(\sqrt{2\gamma}r)}{2 \int_0^1 \Phi(\sqrt{2\gamma}s) ds}, \quad (29)$$

with corresponding energy functionals  $\mathcal{P}_\gamma$ , and notice that the two optimisation problems

$$\mathcal{P} \rightarrow \max \text{ on } \mathcal{S}_\gamma \quad \text{and} \quad \mathcal{P}_\gamma \rightarrow \max \text{ on } \mathcal{S}_{1/2},$$

are equivalent due to  $W \in \mathcal{S}_\gamma \Leftrightarrow W/\sqrt{2\gamma} \in \mathcal{S}_{1/2}$ .

**Example 3.6.** For  $\Phi$  as in (22) the rescaled potentials  $\Phi_\gamma$  from (29) have the complete localisation property on  $[0, 1]$  for  $\gamma \rightarrow \infty$ .

*Proof.*  $\mathcal{P}_\gamma(W_{\text{CL}}) = 1$  holds by construction, and for each  $0 < r_0 < 1$  one can find  $\gamma_0$  such that  $\partial_\gamma \Phi_\gamma(r) < 0$  for all  $0 \leq r \leq r_0$  and  $\gamma \geq \gamma_0$ .  $\square$

More generally, the family  $(\Phi_\gamma)_\gamma$  can be expected to have the complete localisation property for  $\gamma \rightarrow \infty$  provided that  $\Phi$  grows faster than every polynomial. The super-polynomial growth condition is necessary as for every homogeneous potential of degree  $q$  we have  $\Phi_\gamma \equiv \Phi_1$ . This reflects the the homogeneous scaling

$$W \rightsquigarrow \lambda W, \quad \sigma^2 \rightsquigarrow \lambda^{q-2} \sigma^2.$$

and shows that the wave trains for homogeneous or polynomial potentials do not localise in the limit  $\gamma \rightarrow \infty$ .

## 4 Solitons

This section we study soliton solutions to (12), that means we set

$$L = \infty, \quad \mathcal{A} = \bar{\mathcal{A}}, \quad \mathcal{S} = \mathcal{U} \cap \mathcal{N}.$$

Moreover, we assume that the potential energy  $\mathcal{P}$  is genuinely super-quadratic (see Definition 4.4 below) and show that for each  $\gamma > 0$  there exists a maximiser of  $\mathcal{P}$  in  $\mathcal{S}_\gamma = \mathcal{U} \cap \mathcal{N} \cap \mathcal{B}_\gamma$ , which is a soliton according to Theorem 2.3.

In what follows we set

$$P(\gamma) := \sup_{W \in \mathcal{S}_\gamma} \mathcal{P}(W),$$

and in order to compare with the harmonic case we introduce

$$P_{\text{harm}}(\gamma) := \sup_{W \in \mathcal{S}_\gamma} \mathcal{P}_{\text{harm}}(W), \quad (30)$$

where

$$\mathcal{P}_{\text{harm}}(W) = \int_{-L}^L \Phi_{\text{harm}}((\mathcal{A}W)(\varphi)) d\varphi = \frac{1}{2} \beta \| \mathcal{A}W \|_2^2$$

denotes the energy functional corresponding to  $\Phi_{\text{harm}}(r) = \frac{1}{2} \beta r^2$ .

First we show  $P_{\text{harm}}(\gamma) = \beta \gamma$  by studying the maximizing sequence  $(U_n)_n \subset \mathcal{S}_\gamma$  defined by

$$U_n(\varphi) = \begin{cases} \frac{\sqrt{2\gamma}}{\sqrt{n}} \cos\left(\frac{\pi}{2n}\varphi\right) & \text{for } |\varphi| \leq n, \\ 0 & \text{for } |\varphi| \geq n. \end{cases} \quad (31)$$



**Lemma 4.1.** *We have  $\beta\gamma \geq \mathcal{P}_{\text{harm}}(U_n) \geq \beta\gamma(1 - O(n^{-2}))$  for all  $n$ , and hence*

$$P_{\text{harm}} = \sup \mathcal{P}_{\text{harm}}|_{\mathcal{S}_\gamma} = \sup \mathcal{P}_{\text{harm}}|_{\mathcal{B}_\gamma} = \beta\gamma.$$

*Proof.* A direct calculation shows  $\frac{1}{2} \|U_n\|_2^2 = \gamma$ , as well as

$$(\mathcal{A}U_n)(\varphi) = \begin{cases} \Theta\left(\frac{\pi}{4n}\right)U_n(\varphi) & \text{for } |\varphi| \leq n - \frac{1}{2}, \\ 0 & \text{for } |\varphi| \geq n + \frac{1}{2}, \end{cases}$$

and

$$0 \leq (\mathcal{A}U_n)(\varphi) \leq U_n\left(n - \frac{1}{2}\right) = \frac{\sqrt{2\gamma}}{\sqrt{n}} \cos\left(\frac{\pi}{2}\left(1 - \frac{1}{2n}\right)\right) = O\left(n^{-3/2}\right) \quad \text{for } \left|\varphi - n\right| \leq \frac{1}{2}.$$

Moreover, we have

$$\mathcal{P}_{\text{harm}}(U_n) \geq \beta\gamma \Theta\left(\frac{\pi}{4n}\right)^2 \int_{-1+\frac{1}{2n}}^{1-\frac{1}{2n}} \cos\left(\frac{\pi}{2}\varphi\right)^2 d\varphi \geq \beta\gamma(1 - O(n^{-2})),$$

and this implies  $\mathcal{P}_{\text{harm}}(U_n) \rightarrow \beta\gamma$  as  $n \rightarrow \infty$ . Finally, due to Lemma 2.5 we find

$$\mathcal{P}_{\text{harm}}(W) = \frac{1}{2}\beta \int_{\mathbb{R}} |(\mathcal{A}W)(\varphi)|^2 d\varphi = \beta \frac{1}{2} \|\mathcal{A}W\|_2^2 \leq \frac{1}{2}\beta \|W\|_2^2,$$

for all  $W \in \mathcal{B}_\gamma$ , and the proof is complete.  $\square$

**Corollary 4.2.** *We have  $P(\gamma) \geq \beta\gamma = P_{\text{harm}}(\gamma)$  for all  $\gamma \geq 0$ .*

*Proof.* Since the case  $\beta = 0$  is trivial we suppose  $\beta > 0$ . Some elementary analysis shows that the sequence  $(U_n)_n$  from (31) satisfies

$$\mathcal{P}(U_n) \approx \int_{|\varphi| \leq n - \frac{1}{2}} \Phi\left(\Theta\left(\frac{\pi}{4n}\right)U_n(\varphi)\right) d\varphi \approx \frac{1}{2} \int_{|\varphi| \leq n - \frac{1}{2}} U_n(\varphi)^2 (\beta + o(1/\sqrt{n})) d\varphi \approx \beta\gamma,$$

where all approximation errors tend to 0 as  $n \rightarrow \infty$ .  $\square$

As already mentioned in the introduction, solitons are genuinely nonlinear phenomena. In particular, the harmonic chain does not allow for solitons, and thus the supremum in (30) cannot be attained.

**Remark 4.3.** *For  $\beta > 0$  there is no maximiser for  $\mathcal{P}_{\text{harm}}$ .*

*Proof.* Suppose for contradiction the existence of a maximizer  $W_{\text{max}}$  for  $\mathcal{P}_{\text{harm}}$  in  $\mathcal{B}_\gamma$ . Then the Lagrangian multiplier rule implies that there exists a (non-negative) Lagrangian multiplier  $\sigma^2$  such that

$$\sigma^2 W_{\text{max}} = \partial \mathcal{P}[W_{\text{max}}] = \beta \bar{\mathcal{A}}^2 W_{\text{max}},$$

i.e.,  $W_{\text{max}}$  is an  $L^2$ -eigenfunction to  $\bar{\mathcal{A}}^2$  with corresponding eigenvalue  $\sigma^2/\beta$ . This is the desired contradiction because the point spectrum of  $\bar{\mathcal{A}}^2$  is empty due to Lemma 2.5.  $\square$

**On the weak convergence of unimodal, even and non-negative functions in  $L^2(\mathbb{R})$**  For the sake of clarity we proceed with some remarks on the strong compactness of weak convergent sequences from  $\mathcal{U} \cap \mathcal{N} \cap L^2$  as this problem becomes relevant in our existence proof for solitons. Strong compactness criterions are, in principle, provided by the concentration compactness method from [Lio84], but since here we consider only functions from  $\mathcal{U} \cap \mathcal{N}$  the arguments simplify a lot.

Consider a sequence  $(W_n)_n \subset \mathcal{S}_\gamma = \mathcal{U} \cap \mathcal{N} \cap \mathcal{B}_\gamma$  that converges weakly in  $L^2$  to some limit  $W_\infty \in \mathcal{S}_\gamma$ . Passing to a subsequence we can always assume that  $\gamma_n = \frac{1}{2} \|W_n\|_2^2 \rightarrow \bar{\gamma}_\infty$  for some  $\bar{\gamma}_\infty$  with  $\gamma_\infty \leq \bar{\gamma}_\infty \leq \gamma$ , where  $\gamma_\infty = \frac{1}{2} \|W_\infty\|_2^2$ . For  $\gamma_\infty = \bar{\gamma}_\infty$  the convergence of norms enforces the convergence  $W_n \rightarrow W_\infty$  to be strong in  $L^2$ , and we are done in this case.

In the case  $\bar{\gamma}_\infty > \gamma_\infty$  the convergence cannot be strong as some amount of the 'mass' of the measures  $\mu_n = W_n(\varphi)^2 d\varphi$  disappears when passing to  $\mu_\infty = W_\infty(\varphi)^2 d\varphi$ . However, since all functions are non-negative, unimodal, and even, the annihilation of mass is governed by only two elementary processes, compare Figure 7. The weakly convergent sequence can form a *peak at the origin*, and/or a *'fat' tail*, where 'fat' means

$$\lim_{L \rightarrow \infty} \lim_{n \rightarrow \infty} \int_{-L}^L W_n^2(\varphi) d\varphi + \int_L^\infty W_n^2(\varphi) d\varphi > 0,$$

so that some non-negligible amount of the norm is transferred to infinity.




Figure 7: On the weak convergence in  $\mathcal{U} \cap \mathcal{N} \cap \mathcal{B}_\gamma$ . A weak convergent sequence may fail to converge strongly due to the formation of a peak at the origin, and/or a 'fat' tail.

The first observation is that a peak does not contribute to the potential energy  $\mathcal{P}$ . In fact, if the height of the peak is of order  $1/\varepsilon$  with  $\varepsilon \ll 1$ , then the norm constraint  $\frac{1}{2} \|W\|_2 \leq \gamma$  implies that the width of the peak is of order  $\varepsilon^2$ , and thus the peak disappears after applying the averaging operator  $\mathcal{A}$ . More rigorously, Lemma 2.5 guarantees  $\mathcal{A}W_n$  to converge strongly to  $\mathcal{A}W_\infty$  on each compact subset of  $\mathbb{R}$ . Consequently, if the strong convergence fails due to the formation of a peak only, we still have  $\mathcal{P}(W_n) \rightarrow \mathcal{P}(W_\infty)$ .

The formation of a tail, however, is much more crucial as this in general implies  $\mathcal{P}(W_n) \not\rightarrow \mathcal{P}(W_\infty)$ . Remember the maximising sequence  $U_n$  for  $\mathcal{P}_{\text{harm}}$  from (31) having the property that all mass of the measures  $U_n(\varphi)^2 d\varphi$  is contained in a fat 'tail' with increasing support and decreasing height, so that the weak limit is zero. Even worse, since the spectrum of  $\mathcal{A}^2 = \beta^{-1} \partial \mathcal{P}_{\text{harm}}$  is continuous, *each* maximising sequence for  $\mathcal{P}_{\text{harm}}$  is expected to have this property, and we conclude that the formation of tails is directly related to the non-existence of solitons for the harmonic chain.

Our strategy to prove the existence of solitons for nonlinear potentials is to show that each sequence, that maximizes  $\mathcal{P}$  in  $\mathcal{S}_\gamma$ , is *localised*, so a 'fat' tail cannot be formed. To this end we restrict our considerations to the case of *super-quadratic* growth, and derive suitable tightness results under this assumptions.

#### 4.1 Existence of solitons for genuinely super-quadratic $\mathcal{P}$

For the remainder of this section we require super-quadratic growth conditions. We start with the assumptions concerning the energy functional  $\mathcal{P}$  as they appear naturally in our existence proof. Below in §4.2 and §4.3 we then discuss the corresponding properties of the atomic interaction potential  $\Phi$ .

**Definition 4.4.** *The functional  $\mathcal{P}$  is called super-quadratic on  $\mathcal{S}_\gamma$  with  $\gamma > 0$  if*

$$\mathcal{P}(sW) \geq s^2\mathcal{P}(W) \quad \text{for all } W \in \mathcal{S}_\gamma \text{ and } 1 \leq s \leq \frac{\sqrt{2\gamma}}{\|W\|_2}. \quad (32)$$

Moreover,  $\mathcal{P}$  is called genuinely super-quadratic on  $\mathcal{S}_\gamma$  if in addition

$$P(\gamma) > P_{\text{harm}}(\gamma) = \beta\gamma.$$

Notice that for each  $\gamma > 0$  the harmonic functional  $\mathcal{P}_{\text{harm}}$  is super-quadratic, but not genuinely super-quadratic in  $\mathcal{S}_\gamma$ .

**Remark 4.5.** *Let  $\mathcal{P}$  be super-quadratic on  $\mathcal{S}_\gamma$ . Then it is super-quadratic on  $\mathcal{S}_{\tilde{\gamma}}$  for all  $0 \leq \tilde{\gamma} \leq \gamma$ , and we have*

$$\frac{P(\gamma_2)}{\gamma_2} \geq \frac{P(\gamma_1)}{\gamma_1} \geq \beta$$

for all  $0 \leq \gamma_1 \leq \gamma_2 \leq \gamma$ . In particular, if  $\mathcal{P}$  is genuinely super-quadratic for  $\gamma_1 \leq \gamma$ , so it is for every  $\gamma_2$  with  $\gamma_1 \leq \gamma_2 \leq \gamma$ .

**Remark 4.6.** *Let  $\mathcal{P}$  be a super-quadratic on  $\mathcal{S}_\gamma$ . Then, we have*

$$\langle \partial\mathcal{P}[W], W \rangle \geq 2\mathcal{P}(W) \quad (33)$$

for all  $W \in \mathcal{S}_\gamma$ .

*Proof.* In view of the continuity properties of  $\mathcal{P}$  and  $\partial\mathcal{P}$  it is sufficient to consider the case  $\|W\|_2 < \gamma$ . For sufficiently small  $\varepsilon$  (32) implies

$$\varepsilon^{-1}(\mathcal{P}((1+\varepsilon)W) - \mathcal{P}(W)) \geq \varepsilon^{-1}((1+\varepsilon)^2 - 1)\mathcal{P}(W) = (2+\varepsilon)\mathcal{P}(W)$$

and the limit  $\varepsilon \rightarrow 0$  gives (33).  $\square$

We now formulate our main technical result concerning the tightness of maximising sequences. Roughly spoken, ‘fat’ tails are not energetically optimal as their contributions to  $\mathcal{P}$  and  $\mathcal{P}_{\text{harm}}$  are comparable, and peaks are not optimal as they do not contribute to the potential energy at all. These naive explanations can be stated rigorously as follows.

**Lemma 4.7.** *For any  $\delta > 0$  the set*

$$\mathcal{S}_{\gamma,\delta} = \{W \in \mathcal{S}_\gamma : \mathcal{P}(W) - \beta\frac{1}{2}\|W\|_2^2 \geq \delta\}$$

is closed under weak convergence.

*Proof.* Let  $(W_n)_n \subset \mathcal{S}_\gamma$  be a given sequence such that  $W_n \rightarrow W_\infty$  as  $n \rightarrow \infty$  weakly in  $\mathbf{L}^2$ , and  $\mathcal{P}(W_n) \geq \beta\gamma_n + \delta$  with  $\gamma_n = \frac{1}{2}\|W_n\|_2^2$ . Without loss of generality we can assume that  $\frac{1}{2}\|W_n\|_2^2 \rightarrow \bar{\gamma}_\infty$  for some  $\bar{\gamma}_\infty$  with  $\gamma_\infty \leq \bar{\gamma}_\infty \leq \gamma$  and  $\gamma_\infty = \frac{1}{2}\|W_\infty\|_2^2$ . It remains to show that  $\mathcal{P}(W_\infty) \geq \beta\gamma_\infty + \delta$ . For  $n \in \mathbb{N} \cup \{\infty\}$  we set

$$\widetilde{W}_n := W_n|_{[-m,+m]}, \quad \widehat{W}_n := W_n - \widetilde{W}_n, \quad (34)$$

where  $m > 0$  is some constant to be chosen below, and this definition implies

$$\|W_n\|_2^2 = \|\widetilde{W}_n\|_2^2 + \|\widehat{W}_n\|_2^2. \quad (35)$$

Our strategy for this proof is to establish the approximations

$$\mathcal{P}(W_n) \approx \mathcal{P}(\widetilde{W}_n) + \mathcal{P}(\widehat{W}_n), \quad \mathcal{P}(\widehat{W}_n) \approx \mathcal{P}_{\text{harm}}(\widehat{W}_n),$$

where the approximation error becomes arbitrary small if both  $m$  and  $n$  are sufficiently large. To show this we fix  $\varepsilon > 0$ , and suppose  $m$  to be sufficiently large such that

$$\frac{1}{2} \|\widehat{W}_\infty\|_2^2 \leq \varepsilon, \quad \left| \mathcal{P}(W_\infty) - \mathcal{P}(\widetilde{W}_\infty) \right| \leq \varepsilon, \quad \int_{m-1}^m W_\infty(\varphi) \, d\varphi \leq \varepsilon. \quad (36)$$

Such a choice for  $m$  exists as  $\widetilde{W}_\infty \rightarrow W_\infty$  strongly in  $L^2$  as  $m \rightarrow \infty$ . Since  $m$  is finite Lemma 2.5 provides  $\mathcal{A}\widetilde{W}_n \rightarrow \mathcal{A}\widetilde{W}_\infty$  strongly in  $L^2$  as  $n \rightarrow \infty$ , and thus we find

$$\left| \|\widetilde{W}_\infty\|_2^2 - \|\widetilde{W}_n\|_2^2 \right| \leq \varepsilon, \quad \left| \mathcal{P}(\widetilde{W}_\infty) - \mathcal{P}(\widetilde{W}_n) \right| \leq \varepsilon, \quad \left| \int_{m-1}^m W_\infty(\varphi) - W_n(\varphi) \, d\varphi \right| \leq \varepsilon \quad (37)$$

for all sufficiently large  $n$ . Moreover, combining (36)<sub>1</sub> and (37)<sub>1</sub> with (35) and  $\gamma_n \rightarrow \overline{\gamma}_\infty$  shows that

$$\frac{1}{2} \|\widehat{W}_n\|_2^2 = \gamma_n - \frac{1}{2} \|\widetilde{W}_n\|_2^2 \leq \overline{\gamma}_\infty - \frac{1}{2} \|\widetilde{W}_\infty\|_2^2 + C\varepsilon \leq \overline{\gamma}_\infty - \gamma_\infty + C\varepsilon \quad (38)$$

with  $C$  independent of  $n$  and  $\varepsilon$ . In virtue of (36)<sub>3</sub> and (37)<sub>3</sub>, and since all functions  $W_n$  are unimodal, non-negative, and even, we also obtain

$$0 \leq (\mathcal{A}\widehat{W}_n)(\varphi) \leq (\mathcal{A}W_n)(\varphi) \leq (\mathcal{A}W_n)(m - \frac{1}{2}) \leq \int_{m-1}^m W_n(\tilde{\varphi}) \, d\tilde{\varphi} \leq 2\varepsilon$$

for large  $n$  and all  $\varphi$  with  $|\varphi| \geq m - \frac{1}{2}$ . This provides

$$\|\mathcal{A}\widehat{W}_n\|_\infty \leq C\varepsilon, \quad \int_{|\varphi-m| \leq \frac{1}{2}} \Phi(\mathcal{A}W_n) \, d\varphi \leq C\varepsilon, \quad (39)$$

and exploiting the expansions of  $\Phi(r)$  for small  $r$ , compare (14), we find

$$\mathcal{P}(\widehat{W}_n) \leq \frac{1}{2}(\beta + o(\varepsilon)) \int_{\mathbb{R}} (\mathcal{A}\widehat{W}_n)(\varphi)^2 \, d\varphi \leq \frac{1}{2}(\beta + o(\varepsilon)) \|\widehat{W}_n\|_2^2 \leq \beta(\overline{\gamma}_\infty - \gamma_\infty) + C\varepsilon \quad (40)$$

thanks to (38). Finally, due to (34) we have

$$\begin{aligned} \mathcal{P}(W_n) &= \int_{|\varphi| \leq m - \frac{1}{2}} \Phi(\mathcal{A}W_n) \, d\varphi + \int_{|\varphi-m| \leq \frac{1}{2}} \Phi(\mathcal{A}W_n) \, d\varphi + \int_{|\varphi| \geq m + \frac{1}{2}} \Phi(\mathcal{A}W_n) \, d\varphi \\ &= \int_{|\varphi| \leq m - \frac{1}{2}} \Phi(\mathcal{A}\widetilde{W}_n) \, d\varphi + \int_{|\varphi-m| \leq \frac{1}{2}} \Phi(\mathcal{A}W_n) \, d\varphi + \int_{|\varphi| \geq m + \frac{1}{2}} \Phi(\mathcal{A}\widehat{W}_n) \, d\varphi \\ &\leq \mathcal{P}(\widetilde{W}_n) + \int_{|\varphi-m| \leq \frac{1}{2}} \Phi(\mathcal{A}W_n) \, d\varphi + \mathcal{P}(\widehat{W}_n), \end{aligned}$$

and (36)<sub>2</sub> and (37)<sub>2</sub> combined with (39) and (40) imply

$$\mathcal{P}(W_n) \leq \mathcal{P}(\widetilde{W}_\infty) + \mathcal{P}(\widehat{W}_n) + C\varepsilon \leq \mathcal{P}(W_\infty) + \beta(\overline{\gamma}_\infty - \gamma_\infty) + C\varepsilon.$$

Using the assumption we conclude that  $\mathcal{P}(W_\infty) \geq \gamma_\infty + \delta - C\varepsilon$ , and this completes the proof because  $\varepsilon$  was chosen arbitrarily.  $\square$

As a direct consequence of Lemma 4.7 we find in the genuinely super-quadratic case that each maximising sequence must be localized and contains a strongly convergent subsequence.

**Corollary 4.8.** *Let  $\mathcal{P}$  be genuinely super-quadratic on  $\mathcal{S}_\gamma$ , and suppose that the sequence  $(W_n)_n \subset \mathcal{S}_\gamma$  is a maximising sequence for  $\mathcal{P}$  on  $\mathcal{S}_\gamma$ , that means*

$$\lim_{n \rightarrow \infty} \mathcal{P}(W_n) = P(\gamma) = \beta\gamma + \delta > P_{\text{harm}}(\gamma),$$

for some  $\delta > 0$ . Then, there exists a subsequence, still denoted by  $(W_n)_n$ , and  $W_\infty \in \partial\mathcal{S}_\gamma$  such that  $W_n \rightarrow W_\infty$  strongly in  $L^2$ , and hence  $\mathcal{P}(W_\infty) = P(\gamma)$ .

*Proof.* We choose the subsequence and  $W_\infty \in \mathcal{S}_\gamma$  such that  $W_n \rightarrow W_\infty$  weakly in  $L^2$ . Thanks to (32) we know that  $\mathcal{P}(sW_n) \geq s^2\mathcal{P}(W_n)$  for all  $s > 1$ , and this implies  $\gamma_n = \frac{1}{2} \|W_n\|_2^2 \rightarrow \gamma$  as  $n \rightarrow \infty$ , because otherwise the sequence  $(W_n)_n$  could not be maximising. Therefore, with

$$\widetilde{W}_n = \frac{\sqrt{2\gamma}}{\sqrt{2\gamma_n}} W_n \in \partial\mathcal{S}_\gamma$$

we find  $\mathcal{P}(\widetilde{W}_n) \rightarrow \beta\gamma + \delta$  and  $\widetilde{W}_n \rightarrow W_\infty$  weakly in  $\mathcal{S}_\gamma$ , and Lemma 4.7 provides

$$\mathcal{P}(W_\infty) \geq \beta\gamma_\infty + \delta$$

with  $\gamma_\infty = \frac{1}{2} \|W_\infty\|_2^2 \leq \gamma$ . In order to show  $W_\infty \in \partial\mathcal{S}_\gamma$  we use again (32) to obtain

$$\beta\gamma + \delta = P(\gamma) \geq \mathcal{P}\left(\sqrt{\frac{\gamma}{\gamma_\infty}} W_\infty\right) \geq \frac{\gamma}{\gamma_\infty} \mathcal{P}(W_\infty) \geq \frac{\gamma}{\gamma_\infty} (\beta\gamma_\infty + \delta) \geq \beta\gamma + \frac{\gamma}{\gamma_\infty} \delta.$$

Since  $\delta > 0$  we conclude that  $\gamma_\infty = \gamma$ , that means  $\|W_n\|_2 \rightarrow \|W_\infty\|_2$ , and this implies that the convergence  $W_n \rightarrow W_\infty$  is strong in  $L^2$ .  $\square$

The combination of Theorem 2.3 and Corollary 4.8 immediately provides the desired existence result for non-negative and unimodal solitons.

**Corollary 4.9.** *If  $\mathcal{P}$  is genuinely super-quadratic on  $\mathcal{S}_\gamma$ , then there exists a maximiser  $W$  of  $\mathcal{P}$  in  $\mathcal{S}_\gamma$ . This maximiser is a non-negative and unimodal soliton with  $\frac{1}{2} \|W\|_2^2 = \gamma$ .*

Finally, we characterise the soliton speed of maximizers of  $\mathcal{P}$ , and mention that an analogous result holds for wave trains (provided that  $\mathcal{P}$  is super-quadratic).

**Remark 4.10.** *The soliton from Corollary 4.9 is super-sonic, that means  $\sigma^2 > \beta$ .*

*Proof.* Testing the soliton equation (12) with  $W$ , and using (33), we find  $\sigma^2 \|W\|_2^2 \geq 2\mathcal{P}(W) > 2\beta\gamma$ .  $\square$

## 4.2 Criteria for super-quadratic $\mathcal{P}$

**Definition 4.11.** *The potential  $\Phi$  is called super-quadratic on the interval  $[0, \sqrt{2\gamma}]$  if*

$$\Phi(sr) \geq s^2\Phi(r)$$

holds for all  $r \geq 0$  and all  $s \geq 1$  with  $rs \in [0, \sqrt{2\gamma}]$ .

**Remark 4.12.** *If the potential  $\Phi$  is super-quadratic on the interval  $[0, \sqrt{2\gamma}]$ , then  $\mathcal{P}$  is super-quadratic on  $\mathcal{S}_\gamma$ .*

*Proof.* Let  $W \in \mathcal{S}_\gamma$  be fixed, and  $s$  be arbitrary with  $1 \leq s \leq \sqrt{2\gamma}/\|W\|_2$ . For all  $r = (\mathcal{A}W)(\varphi) \leq \sqrt{\|W\|_2^2}$  we have  $rs \leq \sqrt{2\gamma}$ , and hence  $\Phi(s(\mathcal{A}W)(\varphi)) \geq s^2\Phi((\mathcal{A}W)(\varphi))$ , for (almost) all  $\varphi \in \mathbb{R}$ . Finally, integration w.r.t.  $\varphi$  yields the desired result.  $\square$

Definition 4.11 implies that  $\Phi$  is super-quadratic on the interval  $[0, \sqrt{2\gamma}]$  if and only if the function

$$\beta_{\Phi}(r) := 2 \frac{\Phi(r)}{r^2} \quad (41)$$

is non-decreasing on  $[0, \sqrt{2\gamma}]$  (notice that  $\beta_{\Phi}(0) = \beta$  in the sense of Assumption 2.4). To obtain further characterisations of super-quadratic growth we consider the following differential inequalities.

$$(C1) \quad \Phi'(r)r - 2\Phi(r) \geq 0 \text{ for all } r \in [0, \sqrt{2\gamma}],$$

$$(C2) \quad \Phi''(r)r - \Phi'(r) \geq 0 \text{ for all } r \in [0, \sqrt{2\gamma}],$$

$$(C3) \quad \Phi'''(r) \geq 0 \text{ for all } r \in [0, \sqrt{2\gamma}].$$

**Remark 4.13.** For all  $\gamma > 0$  and all sufficiently smooth potentials  $\Phi$  with  $\Phi(0) = \Phi'(0) = 0$  we have

$$(C3) \implies (C2) \implies (C1),$$

where (C1) holds if and only if  $\Phi$  is super-quadratic on  $[0, \sqrt{2\gamma}]$ .

*Proof.* (C3) is equivalent to  $\frac{d}{dr}(\Phi''(r)r - \Phi'(r)) \geq 0$ , and in view of  $\Phi''(0)r - \Phi'(0) = 0$  we conclude that (C3)  $\implies$  (C2). Moreover, the implication (C2)  $\implies$  (C1) can be proven similarly, and (C1) is equivalent to  $\frac{d}{dr}\beta_{\Phi}(r) \geq 0$ .  $\square$

We proceed with a remark concerning the relation between super-quadratic growth and the convexity of  $\Phi$ . In our context, of course, super-quadratic potentials being not convex are forbidden, but solitons can still be shown to exist for such potentials, see [FW94] and our comments below.

**Remark 4.14.** (C2) implies the convexity of  $\Phi$  (on the interval  $[0, \sqrt{2\gamma}]$ ), but there exists non-convex potentials satisfying (C1)

*Proof.* Suppose that  $\Phi$  satisfies (C2). Then, the comparison principle for ODEs gives  $\Phi'(r) \geq \beta r$  for some  $\beta \geq 0$ , and (C2) implies  $\Phi''(r) \geq \beta$  for all  $r > 0$ . Now let  $\eta > 0$  be arbitrary, and consider the potential

$$\Phi_{\eta}(r) = r^2(1 + 2\pi^{-1} \arctan(\eta(r-1))),$$

which is super-quadratic on  $[0, \infty)$  as the function  $\beta_{\Phi_{\eta}}$  is strictly increasing by construction. A direct calculation yields

$$\Phi_{\eta}''(1 + \eta^{-1}) = \pi^{-1}(3 + 3\pi + 2\eta - \eta^2),$$

hence  $\Phi_{\eta}$  is not convex for large  $\eta$ .  $\square$

Another remark concerns the convexity of forces, which become important in the context of atomistic Riemann problems in FPU chains, compare [HR08b].

**Remark 4.15.** (C3) implies the convexity of  $\Phi'$  (on the interval  $[0, \sqrt{2\gamma}]$ ), but there exists potentials  $\Phi$  that satisfy (C2) with non-convex derivative.

*Proof.* The first statement is obvious, and towards the second claim we argue as follows. We choose a non-negative, but not monotonically increasing function  $h$  with  $h(0) = 0$ , and compute  $\Phi'$  as solution to the ODE  $h(r) = \Phi''(r)r - \Phi'(r)$ . Then, each local extremum of  $h$  for  $r > 0$  is a turning point of  $\Phi'$ , and vice versa.  $\square$

### 4.3 Criteria for genuinely super-quadratic $\mathcal{P}$

In order to complete the existence proof for solitons we must show that for a given super-quadratic potential  $\Phi$  the corresponding energy functional  $\mathcal{P}$  is in fact genuinely super-quadratic on  $\mathcal{S}_\gamma$ . In the simplest case there is no harmonic contribution to  $\mathcal{P}$  at all, and then there exist solitons with arbitrary small  $\gamma$ . This holds in particular for all homogenous potentials  $\Phi(r) = cr^\alpha$  with  $c > 0$  and  $\alpha > 2$ .

**Remark 4.16.** *Let  $\Phi$  be super-quadratic on  $[0, \infty)$  with  $\beta = \Phi''(0) = 0$  and  $\Phi(r) > 0$  for all  $r > 0$ . Then,  $\mathcal{P}$  is genuinely super-quadratic on  $\mathcal{S}_\gamma$  for all  $\gamma > 0$ .*

The case  $\beta > 0$  is more involved and needs a better understanding of the balance between the harmonic and non-harmonic contributions to  $\mathcal{P}$ . Our strategy in this case is to find a particular function  $W_0$  such that  $\mathcal{P}(W_0) > \frac{1}{2}\beta \|W_0\|_2^2$ , and this in turn implies the existence of solitons for all  $\gamma \geq \frac{1}{2} \|W_0\|_2^2$ .

**Lemma 4.17.** *Suppose that  $\Phi$  is super-quadratic on  $[0, \infty)$  and that the function  $\beta_\Phi$  from (41) satisfies  $\lim_{r \rightarrow \infty} \beta_\Phi(r) > \frac{3}{2}\beta = \frac{3}{2}\beta_\Phi(0)$ . Then,  $\mathcal{P}$  is genuinely super-quadratic on  $\mathcal{S}_\gamma$  for all sufficiently large  $\gamma$ .*

*Proof.* For fixed  $0 < \varepsilon < 1$  and  $W = \sqrt{2\gamma} W_{\text{CL}}$  with  $W_{\text{CL}}$  as in (23) we find

$$\begin{aligned} \mathcal{P}(W) &= 2 \int_0^1 \Phi(\sqrt{2\gamma}s) ds \geq 2\gamma \int_0^1 \beta_\Phi(\sqrt{2\gamma}s) s^2 ds \\ &\geq 2\gamma\beta_\Phi(0) \int_0^\varepsilon s^2 ds + 2\gamma\beta_\Phi(\varepsilon\sqrt{2\gamma}) \int_\varepsilon^1 s^2 ds = \frac{2}{3}\beta\gamma \left( \varepsilon^3 + \frac{\beta_\Phi(\sqrt{2\gamma}\varepsilon)}{\beta} (1 - \varepsilon^3) \right), \end{aligned}$$

and conclude that  $\mathcal{P}(W) > \beta\gamma$  for all sufficiently large  $\gamma$ .  $\square$

Lemma 4.17 implies the existence of solitons for weakly super-quadratic potentials as for instance

$$\Phi(r) = \frac{\beta}{2}r^2(1 + c \ln(1 + r)), \quad \bar{\Phi}(r) = \frac{\beta}{2}r^2(1 + d \arctan(r))$$

with  $\beta \geq 0$ ,  $c > 0$  arbitrary, and  $d > 0$  sufficiently large.

Next we evaluate the sequence  $(U_n)_n$  from (31) and find an existence criterion for solitons that is very close to that given in [FW94].

**Lemma 4.18.** *Let  $\Phi$  be super-quadratic on  $[0, \infty)$ , and suppose*

$$\Phi(r) \geq \frac{1}{2}\beta r^2 + \varepsilon r^p$$

*for all  $r \geq 0$  with two constants  $\varepsilon > 0$  and  $p > 2$ . Then there exists  $\gamma_0 > 0$  such that  $\mathcal{P}$  is genuinely super-quadratic on  $\mathcal{S}_\gamma$  for all  $\gamma > \gamma_0$ . Moreover,  $\beta = 0$  or  $2 < p < 6$  implies  $\gamma_0 = 0$ .*

*Proof.* Let  $\gamma > 0$  be arbitrary, and consider the sequence  $(U_n)_n$  from (31). Then,

$$\mathcal{P}(U_n) - \mathcal{P}_{\text{harm}}(U_n) \geq \varepsilon \mathcal{P}_{\text{nl}}(U_n)$$

with

$$\begin{aligned} \mathcal{P}_{\text{nl}}(U_n) &= \int_{\mathbb{R}} (\mathcal{A}U_n)^p d\varphi \geq \int_{|\varphi| \leq n - \frac{1}{2}} \left( \Theta\left(\frac{\pi}{4n}\right)U_n \right)^p d\varphi \\ &\geq \left( \Theta\left(\frac{\pi}{4n}\right) \right)^p \left( \frac{2\gamma}{n} \right)^{p/2} n \int_{-1 + \frac{1}{2n}}^{1 - \frac{1}{2n}} \left( \cos\left(\frac{\pi}{2}\varphi\right) \right)^p d\varphi \geq cn^{1-p/2}. \end{aligned}$$

We conclude

$$\mathcal{P}(U_n) \geq \mathcal{P}_{\text{harm}}(U_n) + c\gamma^{p/2}n^{1-p/2} > 0$$

for some positive constant  $c > 0$ , and according to Lemma 4.1 there exists a constant  $\tilde{c} > 0$  such that

$$\mathcal{P}(U_n) \geq c\gamma^{p/2}n^{1-p/2} + \beta\gamma(1 - \tilde{c}n^{-2}).$$

Finally, for  $\beta = 0$ , or  $\gamma$  sufficiently large, or  $-1 + p/2 < 2$  and  $n$  large, we find  $\mathcal{P}(U_n) > \beta\gamma$ .  $\square$

As an application of Lemma 4.18 we find the following existence result for unimodal solitons with non-negative  $W$ : Let  $\Phi(r)$  be analytic with non-negative coefficients, i.e.,  $\Phi(r) = \frac{1}{2}\beta r^2 + \sum_{i=3}^{\infty} \kappa_i r^i$  with  $\kappa_i \geq 0$  for all  $i \geq 3$ . Then  $\Phi$  is super-quadratic on  $[0, \infty)$  and genuinely super-quadratic for large  $\gamma$ . Moreover, if at least one of the coefficients  $\kappa_3, \kappa_4$ , and  $\kappa_5$  is positive, then  $\Phi$  is genuinely super-quadratic for all  $\gamma > 0$ .

Moreover, since the traveling wave equation (12) is invariant under the reflection symmetry

$$W \rightsquigarrow -W, \quad \Phi(r) \rightsquigarrow \Phi(-r),$$

we find also existence results for solitons with non-positive  $W$ . For instance, the Toda potential  $\Phi_{\text{Toda}}(r) = e^{-r} + r - 1$  is not super-quadratic for  $r \geq 0$  but the reflected potential  $\tilde{\Phi}_{\text{Toda}}(r) = e^r - r - 1$  has solitons with arbitrary small  $\gamma$ , compare Lemma 4.18. From this we infer that the Toda chain allows for solitons with  $W \leq 0$ .

Finally, we summarise some other super-quadratic growth conditions for  $\Phi$  under which the existence of solitons was proved by other authors.

1.  $\Phi'(r)r > 2\Phi(r)$ : Friesecke and Wattis [FW94] prove the existence of super-sonic solitons with prescribed potential energy  $\mathcal{P} \geq P_0$  above some critical energy  $P_0 \geq 0$ . Moreover,  $\Phi(r) = \frac{1}{2}\beta r^2 + \varepsilon r^p(1 + o(r))$  with  $\varepsilon > 0$  and  $p$  as in Lemma 4.18 implies  $P_0 = 0$ .
2.  $\Phi(r) = \frac{\beta}{2}r^2 + \Phi_{\text{nl}}(r)$  and  $\Phi'_{\text{nl}}(r)r \geq \alpha\Phi_{\text{nl}}(r)$  for all  $r \geq 0$  and some  $\alpha > 2$ : Smets and Willem [SW97] establish the existence of solitons with prescribed super-sonic speed  $\sigma^2 > \beta = \Phi''(0)$
3.  $\Phi'_{\text{nl}}(r)r \geq \alpha\Phi_{\text{nl}}(r)$  for all  $r \geq 0$  and some  $\alpha > 2$ , or  $\Phi''_{\text{nl}}(r)r \geq \tilde{\alpha}\Phi'_{\text{nl}}(r)$  for all  $r \geq 0$  and some  $\tilde{\alpha} > 1$ : Pankov and Pflüger [PP00] prove under both assumptions the existence of a family of solitons parametrised by  $\sigma^2 > \beta = \Phi''(0)$ .
4.  $\Phi'(r) \geq 0$  for all  $r \geq 0$  and  $\liminf_{r \rightarrow \infty} r^{-\alpha}(\Phi'(r)r - \alpha\Phi(r)) > 0$  and some  $\alpha > 2$ : Schwetlick and Zimmer [SZ07] show that for each super-sonic speed  $\sigma^2 > \sigma_{\text{crit}} \geq \beta = \Phi''(0)$  there exists a soliton.

All these existence results imply that the soliton profile  $W$  belongs to  $\mathcal{N}$ , but since they do not require the convexity of  $\Phi$ , they do not provide  $W \in \mathcal{U}$ .

#### 4.4 Solitons as limits of wave trains

It is very natural to ask whether or not wave trains converge to solitons when the periodicity length  $L$  tends to  $\infty$ . In this section we establish such a convergence result for unimodal and non-negative wave trains. To this end we allow for arbitrary values of  $L \in (0, \infty]$ , and write  $\mathcal{A}_L$  for the operator  $\mathcal{A}$  acting on  $\mathbb{L}^2([-L, L])$ . Consequently, we introduce

$$\mathcal{S}_{L,\gamma} := \mathcal{U} \cap \mathcal{N} \cap \mathcal{B}_{L,\gamma}$$

where  $\mathcal{B}_{L,\gamma}$  denotes the ball of radius  $\sqrt{2\gamma}$  in  $\mathbb{L}^2([-L, L])$ , and consider

$$P_L := \sup_{W \in \mathcal{S}_{L,\gamma}} \int_{-L}^L \Phi((\mathcal{A}_L W)(\varphi)) \, d\varphi.$$



Moreover, we define an *embedding* operator  $E_L : \mathcal{S}_{L,\gamma} \rightarrow \mathcal{S}_{\infty,\gamma}$  by

$$(E_L W_L)(\varphi) = \begin{cases} W_L(\varphi) & \text{for } |\varphi| \geq L, \\ 0 & \text{else.} \end{cases}$$

Inspired by the notion of  $\Gamma$ -convergence we show that the energy of each periodic profile can be approximated by localised profiles, and prove that each localised profile can be recovered by periodic profiles.

**Lemma 4.19.** *For each  $\gamma > 0$  and  $L < \infty$  there exists a constant  $C_{L,\gamma}$  of order  $o(\sqrt{\gamma/L})$  such that*

$$\mathcal{P}_{\infty}(E_L W_L) + C_{L,\gamma} \geq \mathcal{P}_L(W_L) \geq \mathcal{P}_{\infty}(E_L W_L) \quad (42)$$

holds for all  $W_L \in \mathcal{S}_{L,\gamma}$ . Moreover, for any  $W_{\infty} \in \mathcal{S}_{\infty,\gamma}$  there exists a family of functions  $(W_L)_{L < \infty}$  such that

$$\mathcal{P}_L(W_L) \xrightarrow{L \rightarrow \infty} \mathcal{P}_{\infty}(W_{\infty}). \quad (43)$$

*Proof.* First let  $W_L \in \mathcal{S}_{L,\gamma}$  be fixed, and notice that

$$(\mathcal{A}_{\infty} E_L W_L)(\varphi) = (E_L \mathcal{A}_L W_L)(\varphi), \quad ||\varphi| - L| \geq \frac{1}{2}, \quad (44)$$

hold by construction. For  $||\varphi| - L| \leq \frac{1}{2}$  we have  $0 \leq W_L(\varphi) \leq W_L(L-1)$  due to  $W_L \in \mathcal{U} \cap \mathcal{N}$ , and hence

$$0 \leq (\mathcal{A}_{\infty} E_L W_L)(\varphi) \leq (E_L \mathcal{A}_L W_L)(\varphi) \leq \int_{\varphi-1/2}^{\varphi+1/2} W_L(\tilde{\varphi}) \, d\tilde{\varphi} = W_L(L-1).$$

Moreover, from  $W_L \in \mathcal{U} \cap \mathcal{N}$  we infer  $(\mathcal{A}_{\infty} E_L W_L)(\varphi) \leq (E_L \mathcal{A}_L W_L)(\varphi)$  and

$$2\gamma \geq \int_{-L+1}^{L-1} W_L(\varphi)^2 \, d\varphi \geq 2(L-1)(W_L(L-1))^2,$$

and therefore

$$0 \leq (\mathcal{A}_{\infty} E_L W_L)(\varphi) \leq (E_L \mathcal{A}_L W_L)(\varphi) \leq \varepsilon, \quad ||\varphi| - L| \leq \frac{1}{2}. \quad (45)$$

with  $\varepsilon = \sqrt{\gamma/(L-1)}$ . The estimate (42) now follows from (44) and (45) via

$$\begin{aligned} 0 &\leq \mathcal{P}_L(W_L) - \mathcal{P}_{\infty}(E_L W_L) \\ &\leq \int_{-\infty}^{\infty} \Phi\left((E_L \mathcal{A}_L W_L)(\varphi)\right) - \Phi\left((\mathcal{A}_{\infty} E_L W_L)(\varphi)\right) \, d\varphi \\ &\leq \int_{||\varphi| - L| \leq 1/2} \Phi\left((E_L \mathcal{A}_L W_L)(\varphi)\right) - \Phi\left((\mathcal{A}_{\infty} E_L W_L)(\varphi)\right) \, d\varphi \leq f(\varepsilon), \end{aligned}$$

with  $f(\varepsilon) = 2\varepsilon \sup_{0 \leq r \leq \varepsilon} \Phi'(r) = o(\varepsilon)$ . Now let  $W_{\infty} \in \mathcal{S}_{\infty,\gamma}$  be fixed, and define  $W_L \in \mathcal{S}_{L,\gamma}$  by

$$(W_L)(\varphi) = W_{\infty}(\varphi) \text{ for } |\varphi| \leq L.$$

Then, in general we have  $W_{\infty}(\varphi) \neq (E_L W_L)(\varphi) = 0$  for  $\varphi > L$  but always  $E_L W_L \rightarrow W_{\infty}$  strongly in  $L^2(\mathbb{R})$  as  $L \rightarrow \infty$ . This implies  $\mathcal{P}_{\infty}(E_L W_L) - \mathcal{P}_{\infty}(W_{\infty}) \rightarrow 0$ , and thanks to (42) we find (43).  $\square$

Lemma 4.19 now provides the convergence of suprema.

**Corollary 4.20.** *We have  $P_L(\gamma) \xrightarrow{L \rightarrow \infty} P_\infty(\gamma)$ .*

*Proof.* For given  $L < \infty$  let  $\overline{W}_L$  be a maximiser of  $\mathcal{P}$  on  $\mathcal{S}_{L,\gamma}$ . Then (42) implies

$$P_L(\gamma) \leq \mathcal{P}_\infty(E_L \overline{W}_L) + o(\sqrt{\gamma/L}) \leq P_\infty(\gamma) + o(\sqrt{\gamma/L}),$$

and hence  $\limsup_{L \rightarrow \infty} P_L(\gamma) \leq P_\infty(\gamma)$ . Moreover, in view of (43) we have

$$\mathcal{P}_\infty(W_\infty) \leq \liminf_{L \rightarrow \infty} P_L(\gamma)$$

for all  $W_\infty$ , and this shows  $P_\infty(\gamma) \leq \liminf_{L \rightarrow \infty} P_L(\gamma)$ .  $\square$

As a further consequence we find that solitons can be constructed as limits of wave trains. More precisely, Corollary 4.20 combined with Corollary 4.8 provides the following convergence result for maximisers.

**Corollary 4.21.** *Let  $\mathcal{P}$  be genuinely super-quadratic on  $\mathcal{S}_\gamma$ , and for each  $L < \infty$  let  $\overline{W}_L$  be a maximiser of  $\mathcal{P}$  in  $\mathcal{S}_{L,\gamma}$ . Then, for any sequence  $(L_n)_n$  with  $L_n \rightarrow \infty$  there exist a subsequence, still denoted by  $L_n$ , and a maximiser  $\overline{W}_\infty \in \mathcal{S}_{\infty,\gamma}$ , such that  $E_{L_n} \overline{W}_{L_n} \rightarrow \overline{W}_\infty$  strongly in  $L^2(\mathbb{R})$ .*

## Acknowledgement

The author would like to thank Jens D.M. Rademacher for several stimulating discussions, and Michael Helmers for providing the example of Remark 4.14.

## References

- [DEFW93] D. B. Duncan, J. C. Eilbeck, H. Feddersen, and J. A. D. Wattis, *Solitons on lattices*, Physica D **68** (1993), 1–11.
- [DH08] W. Dreyer and M. Herrmann, *Numerical experiments on the modulation theory for the nonlinear atomic chain*, Physica D **237** (2008), no. 2, 255–282.
- [DHM06] W. Dreyer, M. Herrmann, and A. Mielke, *Micro-macro transition for the atomic chain via Whitham’s modulation equation*, Nonlinearity **19** (2006), no. 2, 471–500.
- [DHR06] W. Dreyer, M. Herrmann, and J. Rademacher, *Pulses, traveling waves and modulational theory in oscillator chains*, Analysis, Modeling and Simulation of Multiscale Problems (A. Mielke, ed.), Springer, 2006.
- [FM02] G. Friesecke and K. Matthies, *Atomic-scale localization of high-energy solitary waves on lattices*, Physica D **171** (2002), 211–220.
- [FM03] ———, *Geometric solitary waves in a 2D mass spring lattice*, Discrete Contin. Dyn. Syst. Ser. B **3** (2003), 105–114.
- [FP99] G. Friesecke and R.L. Pego, *Solitary waves on FPU lattices. I. Qualitative properties, renormalization and continuum limit*, Nonlinearity **12** (1999), no. 6, 1601–1627.
- [FPU55] E. Fermi, J. Pasta, and S. Ulam, *Studies on nonlinear problems*, Los Alamos Scientific Laboratory Report LA–1940, 1955, reprinted in: D.C. Mattis (editor), The many body problem. World Scientific, 1993.
- [FV99] A.-M. Filip and S. Venakides, *Existence and modulation of traveling waves in particle chains*, Comm. Pure Appl. Math. **51** (1999), no. 6, 693–735.

- [FW94] G. Friesecke and J.A.D. Wattis, *Existence theorem for solitary waves on lattices*, Comm. Math. Phys. **161** (1994), no. 2, 391–418.
- [Her05] M. Herrmann, *Ein Mikro-Makro-Übergang für die nichtlineare atomare Kette mit Temperatur*, Phd thesis, Humboldt-Universität zu Berlin, 2005.
- [Her09] ———, *Periodic travelling waves in convex Klein-Gordon chains*, in preparation, 2009.
- [HLM94] M.H. Hays, C.D. Levermore, and P.D. Miller, *Macroscopic lattice dynamics*, Physica D **79** (1994), no. 1, 1–15.
- [HR08a] M. Herrmann and J. Rademacher, *Heteroclinic travelling waves in convex FPU-type chains*, submitted, preprint available under [arXiv:0812.1712](https://arxiv.org/abs/0812.1712), 2008.
- [HR08b] ———, *Riemann solvers and undercompressive shocks in FPU chains*, submitted, 2008.
- [Ioo00] G. Iooss, *Travelling waves in the Fermi-Pasta-Ulam lattice*, Nonlinearity **13** (2000), 849–866.
- [Lio84] P.-L. Lions, *The concentration-compactness principle in the calculus of variations. The locally compact case. I*, Ann. Inst. H. Poincaré Anal. Non Linéaire **1** (1984), no. 2, 109–145.
- [PP00] A. Pankov and K. Pflüger, *Traveling waves in lattice dynamical systems*, Math. Meth. Appl. Sci. **23** (2000), 1223–1235.
- [SW97] D. Smets and M. Willem, *Solitary waves with prescribed speed on infinite lattices*, J. Funct. Anal. **149** (1997), 266–275.
- [SZ07] H. Schwetlick and J. Zimmer, *Solitary waves for nonconvex FPU lattices*, J. Nonlinear Sci. **17** (2007), no. 1, 1–12.
- [SZ08] ———, *Existence of dynamic phase transitions in a one-dimensional lattice model with piecewise quadratic interaction potential*, submitted, 2008.
- [Tod81] M. Toda, *Theory of nonlinear lattices*, Springer Series in Solid-State Sci., vol. 20, Springer, Berlin, 1981.